

Original Research Article

<https://doi.org/10.20546/ijcmas.2021.1003.027>

Effect of Integrated Nutrient Management, Cycocel and Crop Geometry on Morphological Features and Yield of Cabbage (*Brassica oleracea* L. var. *capitata*) in Bundelkhand Region

Harpal Singh* and Rajdeep Kudesia

Department of Horticulture, Bundelkhand University, Jhansi, (Uttar Pradesh) India

*Corresponding author

ABSTRACT

In the diversified farming system, integrated nutrient management under different crop geometry along with growth regulators is the most appropriate approach to solve various issues related to productivity, quality as well as sustainability in efficient and economic way. Cabbage is well known for its nutritive and health benefits. A field experiments were conducted to study the effect of vermicompost, inorganic fertilizers and growth retardant (cycocel) with different spacing on cabbage carried out in factorial randomized block design at Experimental Farm of Bundelkhand University, Jhansi (U.P.) during *rabi* season 2017-18 and 2018-19. Results indicated that yield and yield contributing components of cabbage cultivar Pusa Drum Head were found to be highest and significant in the treatment where 50% RDF (Inorganic Fertilizer) + 50% Organic (vermicompost) were applied at wider crop geometry (45cm x 45cm spacing) along with spraying of 300 ppm cycocel (growth retardant). The head yield of cabbage was positively correlated with various yield components. Application of vermicompost along with inorganic fertilizers with cycocel treatment reduces the days taken to head maturity. The experimental findings indicated that reduction in the amount of inorganic fertilizer was not detrimental to total yield to cabbage. So 50% doses of recommended fertilizer would be supplemented with vermicompost. It is also evident that integration of vermicompost and inorganic fertilizer at wider spacing under cycocel treatment enhances the productivity of cabbage beside sustaining soil fertility status.

Keywords

Cabbage,
Vermicompost,
Fertilizer, Cycocel,
Spacing, Yield

Article Info

Accepted:
04 February 2021
Available Online:
10 March 2021

Introduction

The significance of balanced nutrient application is well adopted by farmers for increasing crop productivity. In the diversified farming system, integrated nutrient management under different crop spacing along with growth regulators is the most appropriate approach to solve various issues

related to productivity, quality as well as sustainability. Cabbage is well known for its nutritive and health benefits. It has four distinct uses viz. salads, boiling, pickling and sauerkraut. Cabbage is grown during winter as a major vegetable crop and during summer it is taken as a minor crop or intercrop in Uttar Pradesh. This crops being a grossly feeder, is always been in a high requirement

of nutrients for producing best quality and productivity. On the other hand its productivity is very low in spite of showing great potential in the state. In Uttar Pradesh the production of cabbage is around 302.97 thousand MT from an area of 9.06 thousand hectare with the productivity of 33.44 t/ha (National Horticulture Database 2017-18). Therefore, keeping in view, the field experiments were conducted in Bundelkhand region to study the effect of vermicompost, inorganic fertilizers and growth retardant (cycocel) with different spacing on cabbage.

Materials and Methods

The present study was conducted at Experimental Farm of Bundelkhand University, Jhansi (U.P.) during *rabi* season 2017-18 and 2018-19, Geographically, the research farm of the university is located at 25.45°N latitude and 78.61°E longitude at an altitude of 285 m from mean sea level having a sub-tropical climate. The Investigation was laid out in factorial randomized block design (F.R.B.D) with 27 treatment combinations and three replications. The experiment comprised of three different factors such as 1. Cycocel *viz.* C₁ (0 ppm), C₂ (300 ppm) and C₃ (600 ppm) 2. Fertilizer application *viz.* F₁ (100% RDF Inorganic Fertilizer 120:80:60 NPK kg/ha), F₂ (75% RDF Inorganic Fertilizer + 25% Organic Vermicompost) and F₃ (50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost)) 3. Plant spacing *viz.* S₁ (30 x 30 cm), S₂ (45 x 30 cm) and S₃ (45 x 45 cm). The observation recorded on plant height, number of open leaves, days taken to head maturity, yield and yield attributes such as weight of diameter of head and head yield. All the agronomic were followed as recommended to the crop. One month old cabbage seedlings of Pusa Drum Head were transplanted in the main field at different spacing. Biometrical observations of five randomly samples were taken from each

treatment with standard procedures and techniques. Data recorded were analyzed with the help of statistical programme- STPR- 3 developed by Department of Mathematics and Statistics, College of Basic Science and Humanities, GBPUAT, Pantnagar, Uttarakhand, India.

Results and Discussion

Two years pooled data shown in Table-1 revealed that maximum plant height at 60 days was found maximum (48.19 cm) in treatment C₁F₃S₃ which was statistically at par with C₁F₃S₂. The finding indicated that application of 50 % vermicompost in combinations with 50% inorganic fertilizers at wider spacing significantly affected the plant height of cabbage. This effect might be attributed to the solubilisation of plant nutrients by addition of vermicompost leading to subsequent uptake of NPK by plant. In wider spaced crop proliferous root system developed under above circumstances resulting better absorption of water and nutrients with improved physical environment of soil. The findings are in accordance with the results of Supe and Marbhal (2008), Sharma *et al.*, (2009) and Kumar and Dhar, (2010). Plant height decreased slightly with increasing the cycocel concentration. Plants treated with cycocel were slightly shorter than the controlled plants (C₁). This might be due to the cycocel affected the transport of carbohydrate to the roots *via* decreasing plant height.

The findings are in similar lines with Leclere *et al.*, (2006) while the maximum numbers of open leaves (19.40) per plants were recorded in C₂F₃S₃ which was non-significant with C₂F₃S₂. The response of 50% vermicompost in combination with 50% inorganic fertilizers was found to be higher in number of leaves at wider spaced crop (45x45 cm and 45x30 cm) as compare to the least spaced crop (30x30

cm). This might be due to integration of organic and inorganic fertilizer which might have appeared complementary and supplementary regarded to each other and exhibited into continuous supply of nutrients. Vermicompost imparts favourable soil structure for better root growth, which ultimately increase number of leaves. The higher number of open leaves might also be increased slightly by the reduction in internodal length which might be affected by cycocel treatments. These results are in similar line with the results of Reid and Carr (1967), Hangarge (1997), Sharma and Bhalla (1998) and Meena and Paliwal (2003).

The different level of integrated nutrient management and crop geometry brought out significant response in thickness of main stem at 60 days after transplanting (DAT). The maximum thickness of main stem were recorded in treatment $C_2F_3S_3$ (43.27 mm) which was statistically at par with $C_3F_3S_3$ while minimum was observed in $C_3F_1S_1$ (31.64 mm). The development of plant is reflected through the increase in thickness of main stem which was caused because of the ceaseless accessibility of supplements. The expansion in stem thickness with vermicompost and inorganic fertilizers might be because of more number of leaves and total leaf surface, which lead to increment in photosynthesis.

The use of vermicompost in combination with inorganic fertilizers might have considerably improved the availability of macro and micro nutrients in the soil, as result of which stem thickness has been boosted. In addition to the inhibitory effect of higher dose of cycocel on cell elongation the stem might grown horizontally which ultimately increase the thickness of main stem. These results were in the line of findings reported by Chinaswamy and Marikulandi (1996), Subham (1998), Mehandran and Kumar (1998) and Lendve *et al.*, (2011) in cabbage. With regard to plant

spread the maximum plant spread were recorded in treatment $C_2F_3S_2$ (71.59 cm) which was statistically at par with $C_3F_3S_2$. Use of vermicompost with inorganic fertilizers (RDF) improved the uptake of essential elements. It is because of proliferous root framework created under wide crop geometry which provides better assimilation of water and supplements with improved actual conditions of soil amended by 50% vermicompost along-with 50% inorganic fertilizers.

This impact could be ascribed to the solubilisation of nutrients which accelerate the photosynthetic activity and secretion of growth promoting substances viz., IAA, kinetin, riboflavin and thiamine which result in better plant spread. In addition, application of cycocel also enhances better plant spread which might be due to reduction in vertical growth. Similar results were also recorded by Singh *et al.*, (2004), Choudhary and Choudhary (2005) and Gholampour *et al.*, (2012) in cabbage.

A perusal of data in table 2 revealed that the different level of integrated nutrient management and crop geometry brought out non-significant response in days taken to head initiation. However, the minimum days taken to head initiation was observed in treatment $C_2F_3S_3$ (64.65 days) whereas, maximum days taken to head initiation was recorded in treatment $C_1F_1S_1$ (69.83 days). The decrease in days taken to head initiation with the application of vermicompost and cycocel might be due to verms plays significant role in supply of all essential nutrients.

The results are in accordance with the findings of Ghug *et al.*, (2007), Kumar *et al.*, (2008), Supe and Marbhal (2008) and Velmurugan *et al.*, (2008) in different crops.

Table.1 Effect of integrated nutrient management, cycocel and crop geometry on vegetative characters of cabbage (Two years pooled)

Treatments	Treatment combinations	Plant height (cm) at 60 days	No. of open leaves per plant	Thickness of Main stem (mm)	Plant spread (cm)
C ₁ F ₁ S ₁	0 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 30x30 cm spacing.	41.51	16.90	35.39	64.24
C ₁ F ₁ S ₂	0 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x30 cm spacing.	45.10	17.47	36.05	64.50
C ₁ F ₁ S ₃	0 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x45 cm spacing.	46.08	17.67	35.09	64.44
C ₂ F ₁ S ₁	300 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 30x30 cm spacing.	41.79	17.53	34.27	63.66
C ₂ F ₁ S ₂	300 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x30 cm spacing.	43.73	17.77	34.54	63.77
C ₂ F ₁ S ₃	300 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x45 cm spacing.	45.45	17.87	33.31	63.66
C ₃ F ₁ S ₁	600 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 30x30 cm spacing.	39.13	16.60	31.64	61.22
C ₃ F ₁ S ₂	600 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x30 cm spacing.	42.10	17.07	32.47	62.74
C ₃ F ₁ S ₃	600 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x45 cm spacing.	41.79	17.60	35.79	63.15
C ₁ F ₂ S ₁	0 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 30x30 cm spacing	46.54	17.83	40.42	65.91
C ₁ F ₂ S ₂	0 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x30 cm spacing	46.46	18.07	40.66	66.90
C ₁ F ₂ S ₃	0 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x45 cm spacing	46.54	18.33	41.04	66.45
C ₂ F ₂ S ₁	300 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 30x30 cm spacing	43.85	18.03	40.55	65.78
C ₂ F ₂ S ₂	300 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic	43.85	18.33	40.25	66.32

	(Vermicompost) with 45x30 cm spacing				
C₂F₂S₃	300 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x45 cm spacing	44.94	18.80	41.26	66.12
C₃F₂S₁	600 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 30x30 cm spacing	40.86	17.43	40.62	64.62
C₃F₂S₂	600 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x30 cm spacing	40.86	18.07	38.17	65.25
C₃F₂S₃	600 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x45 cm spacing	40.94	18.77	38.19	68.74
C₁F₃S₁	0 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 30x30 cm spacing	47.17	17.33	41.40	69.28
C₁F₃S₂	0 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x30 cm spacing	48.10	18.30	41.80	70.85
C₁F₃S₃	0 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x45 cm spacing	48.19	18.80	42.57	69.18
C₂F₃S₁	300 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 30x30 cm spacing	43.33	18.40	41.78	69.29
C₂F₃S₂	300 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x30 cm spacing	44.35	18.90	42.56	71.59
C₂F₃S₃	300 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x45 cm spacing	45.44	19.40	43.27	70.20
C₃F₃S₁	600 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 30x30 cm spacing	40.84	18.03	41.50	68.07
C₃F₃S₂	600 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x30 cm spacing	41.92	18.27	41.36	71.18
C₃F₃S₃	600 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x45 cm spacing	43.22	18.57	43.28	70.92
SEm		0.02	0.07	0.05	0.21
CD		0.07	0.21	0.15	0.59

Table.2 Effect of integrated nutrient management, cycocel and crop geometry on days taken to head initiation, head maturity, head size and yield of cabbage (Two years pooled)

Treatments	Treatment combination	Days taken to head initiation (Days)	Days taken to head maturity from planting (Days)	Head size (cm)	Yield (kg/plant)
C ₁ F ₁ S ₁	0 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 30x30 cm spacing.	69.83	85.73	12.11	0.570
C ₁ F ₁ S ₂	0 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x30 cm spacing.	69.53	84.33	12.53	0.950
C ₁ F ₁ S ₃	0 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x45 cm spacing.	69.58	86.47	12.57	1.013
C ₂ F ₁ S ₁	300 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 30x30 cm spacing.	69.16	83.87	13.05	0.953
C ₂ F ₁ S ₂	300 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x30 cm spacing.	68.74	82.73	13.25	1.152
C ₂ F ₁ S ₃	300 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x45 cm spacing.	68.62	82.17	13.27	1.438
C ₃ F ₁ S ₁	600 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 30x30 cm spacing.	69.16	80.40	13.07	0.945
C ₃ F ₁ S ₂	600 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x30 cm spacing.	68.58	81.87	13.33	1.263
C ₃ F ₁ S ₃	600 ppm CCC + 100% RDF Inorganic Fertilizer (120:80:60 NPK kg/ha) with 45x45 cm spacing.	68.53	81.33	13.55	1.550
C ₁ F ₂ S ₁	0 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 30x30 cm spacing	69.41	84.87	12.57	1.525
C ₁ F ₂ S ₂	0 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x30 cm spacing	69.12	83.10	13.09	0.958
C ₁ F ₂ S ₃	0 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x45 cm spacing	69.03	83.43	13.13	1.005
C ₂ F ₂ S ₁	300 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 30x30 cm spacing	68.53	81.73	13.59	1.093
C ₂ F ₂ S ₂	300 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x30 cm spacing	67.83	81.07	13.71	1.382

C₂F₂S₃	300 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x45 cm spacing	67.37	80.60	13.77	1.858
C₃F₂S₁	600 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 30x30 cm spacing	68.08	83.07	13.63	1.402
C₃F₂S₂	600 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x30 cm spacing	64.81	76.60	13.97	1.815
C₃F₂S₃	600 ppm CCC + 75% RDF Inorganic Fertilizer + 25% Organic (Vermicompost) with 45x45 cm spacing	65.78	77.47	14.01	2.117
C₁F₃S₁	0 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 30x30 cm spacing	69.41	85.13	12.83	1.612
C₁F₃S₂	0 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x30 cm spacing	68.74	84.33	13.19	1.368
C₁F₃S₃	0 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x45 cm spacing	68.83	83.43	13.25	1.470
C₂F₃S₁	300 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 30x30 cm spacing	65.35	76.60	14.25	1.068
C₂F₃S₂	300 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x30 cm spacing	65.03	76.10	14.64	1.863
C₂F₃S₃	300 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x45 cm spacing	64.65	75.87	14.72	3.030
C₃F₃S₁	600 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 30x30 cm spacing	65.62	78.50	14.31	1.800
C₃F₃S₂	600 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x30 cm spacing	64.99	76.63	14.68	1.753
C₃F₃S₃	600 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x45 cm spacing	65.07	76.17	14.76	2.733
SEm		0.40	0.37	0.01	0.03
CD		NS	1.04	0.03	0.07

In terms of days taken to head maturity from planting the minimum days taken to head maturity was also observed in treatment C₂F₃S₃ (75.87 days) which was statistically non significant with C₂F₃S₂ and C₃F₃S₃ whereas, maximum days taken to head initiation was recorded in treatment C₁F₁S₃ (86.47 days). In general, the integration of cycocel has significantly reducing the time taken for maturation of head under wide spaced crop nourished with 50% vermicompost + 50% inorganic fertilizers. This might be due to increase in dry matter production during the earlier growth stages because of readily availability of essential nutrient in the soil. The results are in conformity with the findings of Hiwal *et al.*, (2010), Shelke *et al.*, (1999) in brinjal, Shizou *et al.*, (1999) in capsicum and Choudhury and Paswan (2018) reported that the plants treated with cycocel plant growth retardant produced the more compact growth and earlier flower bud initiation in chrysanthemum.

The maximum head size (14.76 cm) was obtained with application of 50% organic manure (vermicompost) + 50% inorganic fertilizer at wider spaced crop (45x45 cm) with higher dose of cycocel (600 ppm) while minimum head size was recorded in C₁F₁S₁ (100% RDF inorganic fertilizers). The increase in head size might be due to the retardation of stem elongation exerted by cycocel which preventing the formation of kaurene, a precursor of gibberellins biosynthesis, subsequently inhibiting or reducing the elongation of stem cells. These results were in the line of findings reported by Gholampour *et al.*, (2012) and Lendve *et al.*, (2011) in cabbage. The data shown in Table-2 expressed that the different level of integrated nutrient management and crop geometry brought out significant response for yield. The maximum significant yield (kg/head) were observed in treatment C₂F₃S₃ (3.030 kg) followed by C₃F₃S₃ (2.733 kg) whereas,

minimum significant yield was recorded in treatment C₁F₁S₁ (0.570 kg).

Application of vermicompost might have considerably increases the availability of macro and micro elements in soil which exerted more yield. The considerable positive increase in yield parameters on account of vermicompost and inorganic fertilizers along with cycocel spray might have resulted to the translocation of nutrients from soil particularly at later stage of crop growth. Application of cycocel increase the growth regulator substance like cytokinein which resulted in enhances the amount of leaf chlorophyll. Then the vegetative growth was efficient to synthesize the improved amount of carbohydrate assimilated by the enhanced rate of photosynthesis. Further increased vegetative growth might have provided more sites for translocation of photosynthesis, which ultimately resulted in increased yield. The plants planted under wider spacing showed more nutrients uptake, light and moisture around compared to plants of narrower spacing, which was possibly the reason of better performance in yield characters and yield of individual cabbage head. These results are in accordance with the findings of Rossini pinto *et al.*, (2005), Dalal *et al.*, (2010) and Chaudhary *et al.*, (2015).

On the basis of present investigation it can be concluded that treatment C₂F₃S₃ (300 ppm CCC + 50% RDF Inorganic Fertilizer + 50% Organic (Vermicompost) with 45x45 cm spacing) performed best in producing higher vegetative growth and yield of cabbage than the other treatments comprised with different levels of cycocel, fertilizer and plant spacing under the present study. It can be further stated that the integrated approach of organic and inorganic fertilization at wider spacing enhances the productivity of cabbage besides sustaining soil fertility.

Acknowledgement

Authors are very grateful to Hon'ble Vice-Chancellor of Bundelkhand University, Jhansi, U.P. and Academic co-ordinator, Institute of Agricultural Sciences, Bundelkhand University campus, Jhansi for providing necessary facilities.

References

- Anonymous., *Indian Horticulture*, Data Base, 2015- NHB. 2017.
- Chaudhary, M., Ajay, K.B. and Parmar, P. Effect of integrated nutrient management on growth, yield attributes and yield of cabbage (*Brassica oleracea* L. Var. *capitata*) under middle Gujarat conditions. *Trends in Biosciences*. 8. 2164-2168. 2015.
- Chinaswamy, K. N. and Mariakulandi, A. Influence of organic and inorganic manure on the firmness and storage life of tomato. *J. South Indian Hort.* 14(1): 36-42. 1996.
- Choudhary, R.K. and Choudhary, D.N. Effect of different levels of N and P on growth, yield and quality of hybrid cabbage. *Haryana Journal of Horticultural Science*. 34(1/2): 145-146. 2005.
- Choudhury, M. T. and Paswan, L. Effect of GA₃ and CCC on growth and flowering of standard chrysanthemum. *International Journal of Agriculture Sciences*.10(11): 6274-6278. 2018.
- Dalal, V.V., Bharadiya, P.S. and Aghav, V.D. Effect of organic and inorganic sources of nitrogen on growth and yield of cabbage (*Brassica oleracea* L. var. *capitata*). *Asian Journal of Horticulture* 5(2): 291-293. 2010
- Gholampur, A., Hashemabadi, D., Sedaghathour, S.H. and Kaviani B. Controlling ornamental cabbage and kale (*Brassica oleracea*L.) growth via cycocel. *Journal of ornamental plants*. 2(2): 103-112. 2012
- Ghug, T.D., Gore, A.K. and Jadhav, S.B. Effect of organic and inorganic nutrient sources on growth, yield and quality of cabbage (*Brassica oleracea*L. var. *Capitata*). *J. Soils & Crops*.17(1):89-92. 2007.
- Hangarge, D.S. Studies on relative efficiency of Vermicompost along with other organic on various soil properties and yield of crops under Chilli-spinach cropping system. M.Sc. (agric.) Thesis. MAU. Prabhani.1997.
- Kumar, A. and Dhar, S. Evaluation of organic and inorganic sources of nutrients in maize (*Zea mays*) and their residual effect on wheat (*Triticum aestivum*) under different fertility levels. *Indian Journal of Agricultural Sciences* 80: 364-71. 2010.
- Kumar, D., Singh, I.P., Singh, B. and Pal, M.K. Effect of integrated nutrient management on growth attributing parameters in cabbage (*Brassica oleracea* L. var. *capitata*). *Progressive Agriculture* 8(2): 243-246. 2008.
- Leclere, M., Caldwell, C.D. and Lade, R.R. Effect of plant growth regulators on propagule formation in *Hemerocallis* spp. and *Hostas* spp. *Hort. Sci.* 47:651-653. 2006.
- Meena, K.K. and Paliwal, R. Growth and yield of cabbage (*Brassica oleracea* L.var. *capitata*) as affected by different nitrogen levels. *Annals of Agricultural Research*. 24(4): 961-963. 2003.
- Mehandran, P.P. and Kumar, N. Effect of organic manure on cabbage cv. Hero (*Brassica oleracea* L. var. *capitata*). *J. South Indian Hort.*45 (5-6): 240-243. 1998.
- Reid, D. M. and Carr, D. J. Effects of a dwarfing compound, CCC, on the production and export of gibberellin-like substances by root systems.

- International journal of agriculture sciences*.11(13):652-654. 1967.
- Rossini, P., Rodrigues, A.C., T.D.J.D., Leits, C.I. and Barbosa, J.C. Growth retardants on development and ornamental quality of potted. 'Liliput' *Zinnia elegans*. *JACQ. Sci. Agric.*62:337-345. 2005.
- Sharma, N.K. and Bhalla, P.C. Influence of integrated nutrient management on growth, yield and economics in okra. *J. Veg. Sci.* 22(1): 1-4. 1998.
- Sharma, A., Kumar, P., Parmar, D.K., Singh, Y. and Sharma, K.C. Bio-inoculants amendment substitutes synthetic fertilizers in cauliflower (*Brassica oleracea* L. var. *capitata*) and influences growth, yield, nutrient uptake and residual soil fertility. *Vegetable Science* 36: 22-26. 2009.
- Shelke, S.R., Adsule, R.N. and Amrutsagar, V.M. Nitrogen management through organics and inorganics in brinjal. *J. of Maha. Agril. Univ.* 24(3):297-298. 1999.
- Shizou, H., Duanwel, Z.H. and Zhu, D.W. Effect of new types of mixed organic manures on yield quality and enzyme activity of chilli and tomato. *J. of Hauzhang Agricultural University.* 18(2): 139-142. 1999.
- Singh, N. P., Bhardwaj, A. K., Kumar A. and Singh, K.M. Modern Technology on Vegetable Production, *International Book Distributing Co.*, New Delhi. 2004.
- Subham, S.S. Effect of organic manures of growth and production of cabbage. *Bull. Pene. Hort.* 16(4):37-41. 1998.
- Supe, V.S. and Marbhal, S.K. Effect of organic manures with graded levels of nitrogen on growth and yield of cabbage (*Brassica oleracea* L.var. *capitata*). *Asian J. Hort.*, 3(1): 48-50. 2008.
- Velmurugan, M., Balakrishnamoorthy, G. and Rajamani, K. Effect of organic manures, biofertilizers and bio-stimulants on growth and yield of cauliflower (*Brassica oleracea* L. var. *botrytis*)cv. Indam 2435. *Crop Res.*, Hisar.35(1/2): 42-45. 2008.

How to cite this article:

Harpal Singh and Rajdeep Kudesia. 2021. Effect of Integrated Nutrient Management, Cycocel and Crop Geometry on Morphological Features and Yield of Cabbage (*Brassica oleracea* L. var. *capitata*) in Bundelkhand Region. *Int.J.Curr.Microbiol.App.Sci.* 10(03): 195-204. doi: <https://doi.org/10.20546/ijcmas.2021.1003.027>