

Original Research Article

<http://dx.doi.org/10.20546/ijcmas.2016.511.027>

A Seasonal Analysis of Micro Zooplankton in the Backwaters of Kottaiappattinam, Tamil Nadu, India

S. Saravanan and R. Sivakami*

P.G and Research Department of Zoology, Arignar Anna Govt. Arts College,
Musiri – 621 211, Tamil Nadu, India

*Corresponding author

ABSTRACT

Keywords

Microzooplankton,
backwater,
Tintinnopsis,
seasons.

Article Info

Accepted:
12 October 2016
Available Online:
10 November 2016

The species composition and diversity of various microzooplankton in the backwaters of Kottaiappattinam, Pudukkottai, Tamil Nadu, India were assessed for three seasons of the year. Pre-summer season recorded 23 species while pre-summer, 38 species and summer 24 species. Thus highest diversity was recorded in the pre-summer season. Among the various groups, tintinnids represented 27 species out the total of 38 species and dominated all the three seasons.

Introduction

Microzooplankton (20-200 μm) form a considerable portion of the zooplankton biomass in marine and estuarine environments (Porter *et al.*, 1985; Pierce and Turner, 1992). Due to their sheer abundance, small size and higher weight specific metabolic rates (Fenchel, 1987; Verity, 1985) microzooplankton play an active role in the transfer of energy to higher trophic levels (Gifford, 1991) from nanoplankton to secondary and tertiary consumers like copepods and fishes (Robertson, 1983; Stoecker and Egloff, 1987).

Unlike the meso and macrozooplankton, the microzooplankton are efficient in consuming

pico and nanoplankton (Nival and Nival, 1976; Johnson *et al.*, 1979; Godhantaraman, 2001) and hence act as a trophic intermediate between pico / nanoplankton and large zooplankton. Further studies show that microzooplankton can be the dominant consumers of phytoplankton production in both oligo and eutrophic regions of the coast and are capable of consuming > 100% of primary production (Elangovan *et al.*, 2012). Hence the present study was aimed at identifying the commonly occurring microzooplankton in Kottaiappattinam backwaters in Tamil Nadu situated in the south-east coast of India.

Materials and Methods

Study area

The present investigation was carried in the backwaters of Kottaipattinam area, Pudukkottai District, Tamil Nadu during the three seasons of the year (Rainy season: October-November 2015; Pre-summer season: January-March, 2016; Summer season: April-June, 2016). Kottaipattinam is situated in Pudukkottai District, Tamil Nadu, with 10.16 Latitude and 78.99 Longitude.

Sample Collection and Analysis

Microzooplankton samples were collected from the surface water by horizontal tow of conical net (0.35 m mouth diameter) made up of bolen silk (mesh size is 54 μm) for twenty minutes. The samples were preserved in 5% neutralized formalin and were used for qualitative analysis. To collect a good number of species and to estimate the abundance of microzooplankton, 100 litre of surface water was filtered with the help of a 10 litre bucket (10 times). As the microzooplankton size is smaller, there are considerable chances of missing few smaller sized forms through the net. Hence, to obtain a reasonable value in the tintinnid abundance, the sedimentation technique was used following the method of Sikhanova (1978). Various biodiversity indices were calculated using the formula of Pielou (1966, 1975 and Gleason (1922).

Results and Discussion

The various microzooplankton of this region belonged to six groups (Tintinnids, Rotifers, Foraminifers, Radiolarians, Ciliates and Copepods) representing a total of 38 species. Among the 38 species, the tintinnids were represented by 27 species while the rotifers

were represented by 5 species, foraminiferans and ciliates by two species each and radiolarians and copepods by one species each (Table-1).

A season wise comparison reveals that a total of 23 species of microzooplankton were recorded during the rainy season. Of these, 14 species belonged to Tintinnids, 4 to rotifers, 2 to foraminiferans and one each to radiolarians, ciliates and copepods. Thus during this season tintinnids formed 60.8% of the microzooplankton while rotifers formed 17.3% and foraminiferans formed 8.6% of the total microzooplankton. On the other hand ciliates, radiolarians and copepods represented only 4.3% each of the microzooplankton.

During the pre-summer season, a total of 38 species of microzooplankton were recorded of which 27 species belonged to tintinnids, 5 species to rotifers, 2 each to foraminiferans and ciliates and one species each to radiolarians and copepods. In terms of percentage, 71% was represented by tintinnids, 13.1% by rotifers, 5.2% each by foraminifers and ciliates and 2.6% each by radiolarians and copepods.

With regard to the summer season, a total of 24 species were recorded of which 19 species belonged to tintinnids, 2 to rotifers and one species each to radiolarians, ciliates and copepods. Surprisingly, foraminifers were absent in this season. In terms of percentage, tintinnids formed 79.1% of microzooplankton followed by rotifers which formed 8.3% and 4.1% each to radiolarians, ciliates and copepods.

An overall percentage composition reveals that tintinnids represented 71% of the total microzooplankton followed by rotifers forming 13.2%. The foraminiferans and ciliate represented 5.2% each followed by

radiolarians and copepods forming 2.6% each. Thus, tintinnids were the most dominating group for all the three seasons. Further, among the three seasons, the most preferred season for microzooplankton appears to be the pre-summer season as the maximum species of microzooplankton were recorded in this season followed by the summer and the rainy season.

A perusal of literature reveals that Prabhu *et al.*, (2005) while analysing the microzooplankton and Senthilkumar *et al.* (2002) from Vellar estuary in Parangipettai coastal waters also recorded maximum diversity during the pre-summer season. They attributed this to the higher productivity of phytoplankton during this season.

Table.1 Shows the Microzooplankton collected from backwaters

S. No.	Species	Rainy Season	Pre-summer Season	Summer Season
Tintinnids				
1.	<i>Tintinnidium primitivum</i>	+	+	+
2.	<i>T. incertum</i>	+	+	+
3.	<i>Tintinnopsis amphora</i>	-	+	-
4.	<i>T. beroider</i>	+	+	+
5.	<i>T. dinecta</i>	-	+	+
6.	<i>T. glans</i>	+	+	-
7.	<i>T. glacialis</i>	-	+	+
8.	<i>T. minuta</i>	+	+	-
9.	<i>T. nucula</i>	+	+	+
10.	<i>T. radix</i>	+	+	+
11.	<i>T. tubulosa</i>	-	+	+
12.	<i>T. butschlii</i>	+	+	+
13.	<i>T. kifoidi</i>	-	+	+
14.	<i>T. nane</i>	+	+	+
15.	<i>T. lohmanni</i>	-	+	+
16.	<i>Codonellopsis orthoceras</i>	+	+	+
17.	<i>C. schabi</i>	-	-	+
18.	<i>Dictyocysta sehaiyai</i>	+	+	+
19.	<i>Stenosemella stenei</i>	+	+	-
20.	<i>Amphorellopsis acuta</i>	-	+	-
21.	<i>Coxliella annulata</i>	+	+	-
22.	<i>Eutintinnus tenuis</i>	-	+	-
23.	<i>Dadayiella bulbosa</i>	+	+	+
24.	<i>Favella brevis</i>	-	+	+
25.	<i>Favella philippinensis</i>	-	+	-
26.	<i>Helicostomella longa</i>	+	+	+
27.	<i>Rhabdonella spiralis</i>	+	+	+

S. No.	Species	Rainy Season	Pre-summer Season	Summer Season
Rotifera				
28.	<i>Brachionus plicatilis</i>	+	+	+
29.	<i>B. argularis</i>	+	+	+
30.	<i>B. urceolaris</i>	+	+	-
31.	<i>Cephalodella gibba</i>	+	+	-
32.	<i>Manostyla bulla</i>	-	+	-
Foraminifera				
33.	<i>Globigerina rubescens</i>	+	+	-
34.	<i>G. plicatilis</i>	+	+	-
Radiolaria				
35.	<i>Acantaria</i> sps.	+	+	+
Ciliata				
36.	<i>Halteria chlorelligera</i>	+	+	+
37.	<i>Lohmaniella spiralis</i>	-	+	-
Copepod nauplius				
38.	<i>Copepod nauplius</i>	+	+	+

Note: ‘+’ represents Presence; ‘-’ represents Absence

Table.2 shows the average diversity index of microzooplankton

S. No.	Details	Rainy Season	Pre-summer Season	Summer Season
1.	Diversity index (H)	2.10	3.62	3.00
2.	Species Evenness (j)	0.56	0.87	0.68
3.	Species Richness (S)	0.94	2.20	1.20

Even though productivity was not analysed in this study, the same reason can be attributed to the maxima obtained as environmental conditions that prevailed would have been more stable when compared to rainy and summer seasons. Literature also reveals that several workers (Mangesh *et al.*, 1996; Krishnamurthy and Naidu, 1977; Qasim and Sengupta, 1981; Prabhu *et al.*, 2005) have also reported

lowest diversity of microzooplankton to occur during the rainy season. This may be attributed to the heavy influx of water from the land into the system as already suggested by a number of workers. In addition, it can also be due to the dispersion of these organisms further away into the sea due to the increased speed of water current as well as winds that are normally associated with this season.

The microzooplankton diversity (H) index (Table 2) in general was found to vary between 2.1 to 3.62 during the three seasons while the evenness (j) ranged from 0.56 to 0.87 and richness from 0.94 to 2.2. While the minimum diversity index was recorded in the rainy season, the maximum was recorded in the pre-summer season.

Among the various groups, tintinnids were found to dominate. Within this group, the species belonging to the genus tintinnids was found to dominate as it was represented by 13 of the 27 tintinnid species. Literature reveals that similar reports were also recorded by Krishnamurthy and Santhanam (1975) and Prabu *et al.*, (2005).

References

- Elangovan, S.S., Kumar, M.A., Karthik, R., Siva Sankar, R., Jayabarathi, R. and Padmavati, G. 2012. Abundance, species composition of microzooplankton from the coastal waters of Port Blair, South Andaman Island. *Aquatic Biosystems*, 8: 20.
- Fenchel, T. 1987. The biology of free-living Phagotrophic Protists. In: *Ecol. Protozoa*. Springer-Verlag: Berlin, p. 197.
- Gifford, D.J. 1991. The protozoan-metazoan trophic link in pelagic ecosystems. *J. Protozoologica*, 38: 81-86.
- Gleason, H.A. 1922. On the relation between species and area. *Ecol.*, 3: 156-162.
- Godhantaraman, N. 2001. Seasonal variations in taxonomic composition, abundance and food web relationship of microzooplankton in estuarine and mangrove waters, Parangipettai region, south-east coast of India. *Indian J. Marine Sci.*, 30: 151-160.
- Johnson, P. W., Sieburth, J. McN, Xu, H-S. 1979. The utilization of crococcoid cyanobacteria by marine protozooplankters but not by calanoid copepods. *Annals de l'Institut Oceanographique*, 58: 297-305.
- Krishnamurthy, K. and Damodara Naidu 1977. Swarming of the tintinnids (Protozoa, Ciliata) in the Vellar Estuary. *Curr. Sci.*, 46: 384.
- Krishnamurthy, K. and Santhanam, R. 1975. Ecology of tintinnids (Protozoa: Ciliata) in Porto Novo region. *Indian J. Mar. Sci.*, 4: 181-184.
- Mangesh, G., Mohanraju, R. and Madhupratap, M. 1996. Studies on microzooplankton from the central and eastern Arabian Sea. *Curr. Sci.*, 71: 874-877.
- Nival, P. and Nival, S. 1976. Particle retention efficiencies of herbivorous copepod (*Acartia clausi* adult and copepodite stages): effect of grazing. *Limnol. Oceanography*, 21: 24-38.
- Pielou, E.C. 1966. The measurement of diversity in different types of biological collections. *J. Theoret. Biol.*, 13: 131-144.
- Pielou, E.C. 1975. *Ecological diversity*. John Wiley and Sons, New York, p. 165.
- Pierce, R.W. and Turner, J.T. 1992. Ecology of planktonic ciliates in marine food webs. Reviews in Aquatic Sciences, Protozoa and their role in marine processes eds. Reid, P. C., Turley, C. M. and Burkill, P. H. NATO ASI Publication, Springer, New York. 6: 139-181.
- Porter, K.G., Sherr, E.G., Sherr, B. F., Pace, M. and Saunders, R. W. 1985. Protozoa in planktonic food webs. *J. Protozool.*, 32: 409-415.
- Prabu, V.A., Perumal, P. and Rajkumar, M. 2005. Diversity and microzooplankton in Parangipettai coastal waters, south-east coast of India. *J. Mar. Biol. Ass. India*, 47: 14-19.
- Qasim, S.Z. and Sengupta, R. 1981.

- Environmental characteristics of the Mandovi-Zuari estuarine system in Goa. *Est. Coast. Shelf Sci.*, 13: 557-578.
- Robertson, J.R. 1983. Predation by estuarine zooplankton on tintinnid ciliates. *Estuarine Coastal and Shelf Sci.*, 25: 581-591.
- Sentthilkumar, S., Santhanam, P. and Perumal, P. 2002. Diversity study in phytoplankton from Vellar Estuary, South-east coast of India. *In: Ayyappan, S., Jena, J. K. and Mohan Joseph, M. Eds), Proc. Fifth Indian Fisheries Foru.* Published by AFR IB, Mangalore and AoA, Bhubaneswar, India, pp. 245-248.
- Stoecker, D.K. and Egloff, D.A. 1987. Predation of *Acartia tonsa* on planktonic ciliates and rotifers. *J. Experimental Marine Biol. Ecol.*, 110: 53-68.
- Verity, P.G. 1985. Grazing, respiration, excretion and growth rates of tintinids. *Limnol. Oceanography*, 30: 1268-1282.

How to cite this article:

Saravanan, S., and Sivakami, R. 2016. A Seasonal Analysis of Micro Zooplankton in the Backwaters of Kottaipattinam, Tamil Nadu, India. *Int.J.Curr.Microbiol.App.Sci.* 5(11): 257-262. doi: <http://dx.doi.org/10.20546/ijcmas.2016.511.027>