

Original Research Article

<https://doi.org/10.20546/ijcmas.2017.610.356>

Prevalence of Candida and Trichomonas in Vaginal Discharge of Rural Females not Seeking Medical Help: Microscopy and its Role in Vaginitis

Rachna Tewari*, Motilal P. Sah and Mridu Dudeja

Department of Microbiology, HIMSR, Jamia Hamdard, New Delhi, India

*Corresponding author

ABSTRACT

In developing countries complaint of abnormal vaginal discharge is the major problem in females of reproductive age group. However most of the rural women do not contact medical facilities in time for adequate treatment due to many social factors or lack of awareness about the disease which leads to complications, risk of increased sexual transmission and insufficient data to deal with. This study is aimed to assess the prevalence of vaginal candidiasis and trichomoniasis among rural females not attending any medical clinic, to raise awareness about its prevention and treatment to reduce its complications. A total of 200 females with complaint of leucorrhoea were selected. Speculum guided high vaginal swabs were collected and processed by KOH mount and Grams stain in field and confirmed by culture on Sabrauds Dextrose agar (SDA) in lab. Out of the 200 females examined 73 (36.5%) had genital infection. Among the positive cases Candida sp. were found in 24 (12%) cases. *Trichomonas vaginalis* were obtained in 9 (4.5%) cases in this study. Prevalence was found significantly higher in lower qualification ($p < 0.01$). Microscopy is simple, cheap and reliable screening method to detect the prevalence of vaginitis in rural areas. The high rate of unnoticed genital infections warrants the need of constant health care and sensitization to prevent complications and acquisition and transmission of sexually transmitted diseases. So, it is essential to detect the disease and treat the cases in time in neglected and knowledge deprived areas.

Keywords

Candida,
Trichomonas,
Vaginal discharge.

Article Info

Accepted:
26 September 2017
Available Online:
10 October 2017

Introduction

Abnormal vaginal discharge with or without other symptoms, is one of the major problem in developing countries. It can be infective or non-infective.

In rural areas it has been noted that this morbidity is tolerated silently by women due to various social factors as poor awareness, gender inequalities, lack of proper education and lack of focused counselling and financial dependence. Candidiasis and Trichomoniasis are two of the most common causes of

vaginal discharge⁽¹⁻⁶⁾. Candida is a yeast like fungus, normally found as the lower genital tract flora in 20-50 % of healthy asymptomatic females⁽⁷⁻⁸⁾. Carrier rates are higher in women using broad spectrum antibiotics⁽⁹⁾, pregnant, diabetic women⁽¹⁰⁻¹¹⁾ and women with HIV/AIDS^(8,12,13). *Candida albicans* is the commonest cause for most cases of vulvovaginal candidiasis (VVC) than other species of Candida. It is characterized by severe itching, burning sensation, redness and curdy white discharge^(9,14). In unattended

pregnant females can lead to preterm labour and low birth weight. Also these females are at risk of acquiring other STDs more.

Trichomoniasis is caused by a protozoa, *Trichomonas vaginalis* which is a sexually transmitted parasite resulting in discomfort, frothy yellow greenish vaginal discharge, irritation, pain in vulva, perineum, dyspareunia, dysurea and psychosocial distress in the infected patients. It has also been reported to cause complication in pregnant women like post abortion infection, post caesarean infection, preterm birth, low birth weight infants and preterm labour^(15,16).

Asymptomatic or mild cases of Candidiasis and Trichomoniasis may not be taken seriously by rural patients and may lead to severe complications like infertility or complications associated with pregnancy and make them vulnerable to other type of sexually transmitted diseases (STDs).

Keeping the above fact in view this study was carried out to see the hidden infection by organizing camps and door to door service to find out affected women who need treatment and counseling for these infections at right time so that spread and complications can be prevented.

Materials and Methods

A cross sectional study was conducted by the Department of Microbiology, HIMSR, New Delhi, in March 2017. Camps were organized in slums and rural areas. A total of 200 women with complaints of vaginal discharge were recruited. After taking consent complete history and general and gynaecological examination was carried out. Three high vaginal sterile swabs from lateral wall and posterior fornix were collected using a speculum during comprehensive pelvic examination. The nature of discharge (amount, colour, character, and smell) was

evaluated. First swab is used to prepare Wet mount to see motility of *Trichomonas vaginalis*, fungal buds, bacteria and pus or other type of cells and to make KOH preparation. Second swab of each patient was used to make smears for Grams and Giemsa stain. Third swab was inoculated in normal saline and carried to microbiology lab at sterile conditions for culture. Blood Agar (BA) and Sabrauds Dextrose agar (SDA), Hi Media, Mumbai was inoculated for swabs showing budding yeast in Grams stain and KOH mount⁽¹⁷⁻¹⁸⁾.

Inclusion criteria

Females of age 18-48 years with complaint of excessive vaginal discharge.

Those who were willing for speculum examination

Exclusion criteria

Not giving consent for speculum examination.

Women who were menstruating or have attained menopause.

Bacterial and non-infective cases were excluded.

Identification of organisms

For Candida

KOH preparation – Little vaginal secretion was taken on clean slide and a drop 10% KOH was mixed and covered with cover slip. Microscopy under 10x and 40x was performed. Round and oval shaped budding yeast cell were identified as Candida.

Grams stain- Gram positive budding yeast cell were identified as Candida in Gram stained vaginal smear.

Culture – It was done on SDA (Sabrauds Dextrose Agar) in lab. Cream coloured, smooth, moist colonies were confirmed as *Candida* by making their Grams Stain.

GTT – Isolated candida colonies were transferred to human sera and seen for tube like yeast cell and pseudohyphae were seen under microscope⁽¹⁸⁾.

For *Trichomonas vaginalis*

Wet mount: motile trophozoites were identified.

Giemsa Stain: oval trophozoites were seen⁽¹⁸⁾.

Ethical clearance was obtained from the department. Eligible females were informed and counselled about the benefits and risk of studies. The consent of all patients were verbally taken before collecting sample and data. Data were analysed using MS excel and appropriate statistical methods.

Results and Discussion

In the current study of two hundred females, with complaint of vaginal discharge following observations were noted. Out of these 200 women 73 (36.5%) were diagnosed to be having infective vaginitis. The prevalence of candidiasis was found in 24 (12%) females out of which majority i.e. 22 (90%) were confirmed as *Candida albicans* and 10% as other *Candida* species. *Trichomonas vaginalis* was identified in 9 (4.5%) females. No case of mixed infection was found (Table 1).

Candida was found in high frequency (52.7%) in age group 39-48 years of females. Prevalence of vaginal candidiasis and Trichomoniasis was found higher in unemployed 25 (20.4%) than employed women 8 (10.1%). No significant correlation

was found with educational status in these areas. Lower incidence 14.8% of infection was found in not educated females where as 18% in higher qualified and 21.6% in mediocre (Table 3).

24 (12%) patients were identified as having infection with *Candida species*. By performing microscopic examination in field 75% strains can be isolated by KOH mount and 83.3% by Grams staining when we compared it with culture all (100%) (Table 3). Germ tube test was found positive for 91.6% strains. *Trichomonas vaginalis* was identified in 9(4.5%) patients by microscopic examination by wet mount and Giemsa Stain (Table 4).

Infection in the genital tract produce serious complications and can be transmitted to others^(19- 20). The doctors or health workers commonly encounter the females associated with vaginal discharge with or without itching or foul smell. Infectious vaginitis primarily caused by bacteria, yeast and protozoa account for 90% of the disease. Vulvovaginal candidiasis and trichomoniasis are the leading cause of vaginitis^(21 -22).

Mostly these studies have been conducted in hospitals in urban area. In our study we are reporting about the females who themselves have not contacted doctors but have been contacted by the health care workers to find out the hidden infection in the community which should also be prevented and treated in time to control the transmission. So this prevalence should be added to the prevalence being reported till now.

In our study we have reported 12 % females having candidiasis which is higher than reported by other studies⁽²³⁻²⁴⁾ which may be due to their large sample size. However it was slightly less as compared to 22% reported by Venugopal *et al.*,⁽²⁵⁾.

Table.1 Etiology of vaginal discharge and its prevalence

Aetiology	Number (Total -200)	Percentage
<i>Candida albicans</i>	17	8.5%
Candida species	07	3.5%
<i>Trichomonas vaginalis</i>	9	4.5%
Other infection	40	20%
No organism found	127	63.5%

Table.2 Demographic characteristics associated with infection

Level of Education	Number (N)	Organism	number (n)	(Percentage) n/Nx100
Less than high school	108	Candida	12	11.1
		Trichomonas	4	3.7
More than high school	72	Candida	9	12.5
		Trichomonas	4	5.5
Not educated	20	Candida	3	15
		Trichomonas	1	6.6
Employed	78	Candida	6	7.6
		Trichomonas	2	2.5
Unemployed	122	Candida	18	14.7
		Trichomonas	7	5.7
AGE in years 18-28	121	Candida	7	5.7
		Trichomonas	3	2.4
29-38	60	Candida	7	11.6
		Trichomonas	4	6.6
39-48	19	Candida	10	52.6
		Trichomonas	2	10.5
Pregnant	28	Candida	9	32.1
		Trichomonas	1	3.5
Not pregnant	172	Candida	15	8.7
		Trichomonas	8	4.6

Table.3 Lab result of *Candida* species

Positive specimen	KOH mount 18/24(75)	Grams stain 20/24 (83.3)	Culture 24/24 (100)	Germ tube 22/24
12	+	+	+	11
8	-	+	+	8
4	+	-	+	3

Table.4 Lab result of *Trichomonas vaginalis*

Positive specimen	Wet mount 7/9	Giemsa staining 7/9
5	+	+
2	-	+
2	+	-

Table.5 Prevalence of candidiasis and *Trichomonas vaginalis* reported earlier and in present study

Author	Year	% age	Place of study	Journal
Swetha Venugopal, Kannan Gopalan	2017	Candida 22% Trichomonas 25%	Salem. Tamil Nadu	Indian Journal of Sexually Transmitted Diseases and AIDS Volume 38
B. Anuradha*, M.C.K. Joanna and M. Praveena	2015	Trichomonas 22.22%	Khammam, India	IJCMAS
R Bansal, N Jindal, B Chand	2013	Trichomonas 6.1%	Punjab India	Ijmm
Alli <i>et al.</i> ,	2011	Candida 22%	Nigeria	JMBR
Madhivanan <i>et al.</i> ,	2008	Candida 30.8% Trichomonas 8.2%	Mysore India	IJMM

Trichomoniasis is a common STD found in humans. We have reported prevalence of 4.5% which is equivalent to reported by other studies. It has been found that prevalence varies from place to place and with social condition from study to study and reported to be low as 2-8%. Whereas high 22-45%.

This may be due to the high risk nature of group ⁽²⁵⁻²⁷⁾. A very high prevalence of TV 47% was reported ⁽²⁸⁾. Studies have indicated

it as reemerging pathogen and found to correlate with age.

Despite the various techniques available and found to be little more sensitive for diagnosing candida species and Trichomoniasis wet mount with clinical features still comes out as reliable screening test (Tables 4 and 5) in rural areas. Although many studies ⁽²⁹⁾ have found more positivity by culture (13.9%) then by wet mount (11.

4%). Microscopy is reliable, inexpensive method, which can be done in field with no requirement of specialist.

Microscopy in the field, substantiated with sign and symptoms is easy, inexpensive and reliable methods to detect the infection in rural females not attending clinics for treatment. It is highly recommended that STDs should be prevented, diagnosed and treated early to avoid complications and reduce transmission. This study was done to create awareness in rural and slum area about health care facilities and self-concern in females for their health. Camps and door to door surveys should be performed to approach these types of unaware females.

References

1. Joshi A, Dhapola M, Pelto PJ. Gynaecological Problems: Perceptions and Treatment-seeking Behaviours of Rural Gujarati Women. In: Michael A Koenig, Shireen Jejeebhoy, John C Cleland and Bela Ganatra (eds.) Reproductive Health in India New Evidence. Jaipur: Rawat Publications. 2008; 133-58.
2. Bhatia, Bhatia JC, Cleland J, Bhagavan L, Rao NS. Levels and determinants of gynecological morbidity in a district of south India. *Stud Fam Plann* 1997; 28:95-103.
3. Gittlesohn J, Holmes KK, Pilot P, Et al (eds.) Listening to Women Talk about Their Health: Issues and Evidence from India. New Delhi: Har-Anand Publications. 1994; 21-36.
4. Oomman, N. Poverty and pathology: Comparing rural Rajasthani women's ethnomedical models with biomedical models of reproductive morbidity: Implications for women's health in India. Dissertation thesis submitted to John Hopkins University, Baltimore, USA. 1996.
5. Wasserheit JN, Harris JR, Chakraborty J, Kay BA, Mason KJ. Reproductive tract infections in a family planning population in rural Bangladesh. *Stud FamPlann* 1989; 20:69-80.
6. Younis N, Khattab H, Zurayk H, el-Mouelhy M, Amin MF, Farag AM. A community study of gynecological and related morbidities in rural Egypt. *Stud FamPlann* 1993; 24:175-86.
7. McClelland RS, Richardson BA, Hassan WM, Graham SM, Kiarie J, Baeten JM, Mandaliya K, Jaoko W, Ndinya-Achola JO, Holmes KK. 2009. *J. Infect. Dis.* 15; 199(12): 1883-1890.
8. Adad SJ, de Lima RV, Sawan ZT, Silva ML, de Souza MA, Saldanha JC, Falco VA, da Cunha AH, Murta EF. 2001. *Sao Paulo Med J.* 119(6):200-205.
9. Singh SI. 2003. Treatment of vulvovaginal candidiasis. *Clin. Rev. CPJ/RPC.* 136(9): 26-30.
10. Donders GG, (2002). *Curr Infect Dis Rep.* 4(6): 536-539.
11. de Leon EM, Jacober SJ, Sobel JD, Foxman B (2002). *BMC Infect Dis.* 2(1). doi:10.1186/1471-2334-2-1.
12. Reed BD, Zazove P, Pierson CL, Gorenflo DW, Horrocks J. 2003. *J. Women's Health (Larchmt).* 12(10): 979-89.
13. Duerr A, Heilig CM, Meikle SF, Cu-Uvin S, Klein RS, Rompalo A, Sobel JD, (2003). *Obstet Gynecol* 101(3): 548-56.
14. Mackay MD. Gynecology. In: Tierney LM, McPhee SJ, Papadakis MA, editors. Current Medical Diagnosis and Treatment. Connecticut: Appleton and Lange; 1998:691-723.
15. Graves A, Gardner WA jr. Pathogenicity of *Trichomonas vaginalis*. *Clin ObstetGynaecol* 1993; 36:145-151.
16. Nanda N, Michel R G, Kurdgelashvili G, Wendel KA. Trichomoniasis and its treatment. *Expert Rev Anti Infect Ther* 2006; 4(1): 125-35.
17. Cheesebrough M. 2006. District

- laboratory practice in tropical countries. Part 2. Cambridge University Press, United Kingdom; p.434
18. Bailey and Scotts Bailey, W. R., Scott, E. G., Finegold, S. M., and Baron, E. J. (1986). *Bailey and Scott's Diagnostic microbiology*. St. Louis, Mosby.
 - 19 .Rajalakshmi R, Kalaivani S. Prevalence of asymptomatic infections in sexually transmitted diseases attendees diagnosed with bacterial vaginosis, vaginal candidiasis, and trichomoniasis. *Indian J Sex Transm Dis* 2016; 37:139-42
 20. Akingbade OA, Akinjinmi AA, Awoderu OB, Okerentugba PO, Okonko IO. Prevalence of *Candida albicans* amongst women attending health centres in Abeokuta, Ogun State, Nigeria. *N Y Sci J* 2013; 6: 53.
 21. Fox KK, Behets FM. Vaginal discharge. How to pinpoint the cause. *Postgrad Med* 1995; 98: 87-90.
 22. Vijaya Mn D, Umashankar K, Sudha, Nagure AG, Kavitha G. Prevalence of the *Trichomonas vaginalis* infection in a tertiary care hospital in rural Bangalore, Southern India. *J Clin Diagn Res* 2013; 7:1401-3.
 23. Pawanarkar J, Chopra K. Prevalence of lower reproductive tract infection in infertile women. *Health Popul Perspect Issues*. 2004; 27: 75.
 24. Gupta V, Chatterjee B, Prasad D. Clinical spectrum and microbial etiology of reproductive tract infections in rural women in the hills of North India. *J Obst Gynaecol India* 2002; 52:130-4.
 25. Venugopal S, Gopalan K, Devi A, Kavitha A. Epidemiology and clinico-investigative study of organisms causing vaginal discharge. *Indian J Sex Transm Dis* 2017; 38: 69-75
 26. Bachmann LH, Hobbs MM, Seña AC, Sobel JD, Schwebke JR, Krieger JN, *et al.* *Trichomonas vaginalis* genital infections: progress and challenges. *Clin Infect Dis* 2011; 53: S160-72.
 27. Lally P, van Jaarsveld CH, Potts HW, Wardle J. How are habits formed: Modelling habit formation in the real world. *Eur J Soc Psychol* 2010; 40:998-1009.
 28. Manson PR, Super H, Fripp PJ. Comparison of four techniques for the routine diagnosis of *Trichomonas vaginalis* infection. *J Clin Pathol* 1976; 29: 154-7.
 29. de Lemos PA, Gracia-Zapata MT, Guimaraes NM, Moris RG. Comparison of methods for identification of *Trichomonas vaginalis* in HIV positive and negative women. *Int J Trop Med* 2009; 4:76-81.

How to cite this article:

Rachna Tewari, Motilal P. Sah and Mridu Dudeja. 2017. Prevalence of *Candida* and *Trichomonas* in Vaginal Discharge of Rural Females not Seeking Medical Help: Microscopy and Its Role in Vaginitis. *Int.J.Curr.Microbiol.App.Sci*. 6(10): 3023-3029.
doi: <https://doi.org/10.20546/ijemas.2017.610.356>