

Original Research Article

<https://doi.org/10.20546/ijcmas.2017.609.096>

Efficacy of Coir Waste Compost and Urea at Different Levels on the Incidence of Sheath Blight of Rice

A. Sudha^{1*}, M. Senthilkumar² and M. Vijayakumar³

¹Department of Plant Pathology, Krishi Vigyan Kendra, Tamil Nadu Agricultural University, Coimbatore, Salem district 636 203, Tamil Nadu, India

²Department Agricultural Entomology, KVK Salem, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India

³Department of Agronomy, KVK Salem, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India

*Corresponding author

ABSTRACT

Keywords

L. connatus,
P. sajor- caju,
Coir waste, Urea,
Phosphorus,
Potassium,
Rhizoctonia solani.

Article Info

Accepted:
14 July 2017
Available Online:
10 September 2017

Lentinus connatus treated coir waste and urea at all levels significantly reduced the intensity of sheath blight, when compared with *Plerotus sajor-caju*. The treatments coir waste compost (*L. connatus*) @ 18.5 t + urea @ 60 kg N/ha and composted coir waste (*L. connatus*) @ 15.5 t/ha + urea @ 60 kg N / ha were found superior and significantly reduced the intensity of sheath blight (Gr. 3.00 and 3.52), compared to the existing fertilizer recommendation of 12.5t composted + 120 kg N/ha, where maximum disease intensity of Gr. 9.75 (sheath blight). The similar trend was also observed with *P. sajor - caju*. The above said best treatments also significantly reduced the disease intensity of sheath blight (Gr. 5.60 and 6.20). The earlier treatments not significantly reduced the disease intensity and on par with existing recommendation.

Introduction

In India, vast quantities of renewable agro and agro-industrial wastes are generated every year (Madan *et al.*, 1987). More than half of the total produce from the land remains unused as wastes (Zadrazil, 1984). The abundant availability of these crop residues, the problems associated with its disposal, the associated environmental pollution problem and the physio-chemical characteristics of these materials attracted the attention of

technologists in finding productive use of these crop residues.

The current boom in inorganic fertilizer prices and low availability of organic manures, limit the use of these viable source of nutrients in agriculture.

There have been many reports of studies on the bioconversion of lignocellulosic materials

into a potential biomanure (Anderson *et al.*, 1971; Zadrazil, 1975; Beg *et al.*, 1986).

Efficient degradation of complex lignocellulosic wastes by *Pleurotus* spp. largely depends upon the synergistic action of extracellular enzymes like cellulases and lignases. These enzymes have been reported to narrow down the C: N ratio in various organic residues, indicating the better rate of lignocellulosic degradation (Sheeba *et al.*, 1999; Singh and Griensven, 2000). Hence, soil application of such a degraded crop residues improves the nutritional status of the soil and suppresses soil borne pathogens (Rajan, 1980; Sheela, 1992).

In a search for an efficient biodegrader for the production of high quality biomanure, the present study was undertaken to probe the potentially of a new mushroom fungus viz., *Lentinus connatus* Berk, isolated from the dead woods of cashew (Lakshmanan, 2002), on comparison with existing biodegrader.

Materials and Methods

Preparation of substrates for composting

The substrates viz., paddy straw, sugarcane trash and sugarcane bagasse were chopped off 5 to 10 cm length. The substrate coir waste was used without chopping. They were solar dried for 5 - 6h. Enough water was added to make it sufficiently moist (40 - 50%). The moisture held in substrate was calculated by gravity method (Piper, 1966).

Method of composting

Heap method of composting described by Manickam *et al.*, 1980) was followed. For composting 20 kg of moist substrate, 20 g fungal culture (*L. connatus* / *P. sajor caju*) and 100 g urea were used. Two kg of substrate was spread uniformly (75 x 45 cm)

in a shaded place and 4 g of fungal culture (*L. connatus* / *P. sajor caju*) was sprinkled over it. This was to be alternated with 2 kg of substrate and 20 g of urea. The alternate layers of substrate + culture and substrate + urea were stacked upto the height of 1m. Substrate only with *L. connatus* / *P. sajor-caju* was also maintained for comparison. The heap was allowed to degrade for a month. For each substrate / commonly grown rice cultivar ADT - 36, was sown in peat trays (10x20x40 cm). Four weeks after germination, the seedlings were transplanted to 10cm dia earthen clay pots filled with 4 kg of clay soil @ 3 seedlings per pot. Before transplanting, 50% of the total inorganic nitrogen and 100% of the total P₂O₅, K₂O and coir waste composted by *L. connatus* / *P. sajor-caju*, were applied at basal separately.

On 45th and 60th day after transplanting, the remaining 50% of total inorganic nitrogen was applied in two split doses. Three replicated pots for each fertilizer dose / composted coir waste (*L. connatus* / *P. sajor-caju*), were maintained in a randomized complete block design.

Treatments details

Fertilizer dose (ha)

T1: 12.5 t compost + 120 kg N + 38 kg P₂O₅ + 38 kg K₂O

T2: 12.5 t compost + 90 kg N + 38 kg P₂O₅ + 38 kg K₂O

T3: 12.5 t compost + 60 kg N + 38 kg P₂O₅ + 38 kg K₂O

T4: 15.5 t compost + 90 kg N + 38 kg P₂O₅ + 38 kg K₂O

T5: 15.5 t compost + 60 kg N + 38 kg P₂O₅ + 38 kg K₂O

T6: 18.5 t compost + 90 kg N + 38 kg P₂O₅ + 38 kg K₂O

T7: 18.5 t compost + 60kg N + 38 kg P₂O₅ + 38 kg K₂O

Artificial inoculation of fungal pathogens

Inoculum production

Preparation of rice grain inoculum

A rice grain mixture (cv.ADT 36) consisting of 2/3 filled grains and 1/3 chaffy grains and weighing 30 g in a 100 ml Erlenmeyer flask, after overnight presoaking in fresh water. After autoclave sterilization at 151b pressure for 2 h, the flask was inoculated with a 9 mm disc of 10-day-old PDA culture of *R. solani*. They were incubated at 25 ±3°C for 15 days and then used for inoculation (Amin *et al.*, 1974).

Method of inoculation

Inoculation was done at active tillering phase, for *T. cucumeris*. In case of *T. cucumeris* rice seedlings were artificially inoculated by inserting the single rice grain culture (15 day-old), being placed inside base of the sheath enclosing culm.

Disease intensity

Individual tillers were assessed at 3 day intervals for fungal disease infection on a 0-9 scale, using Standard Evaluation System for Rice Diseases (Anon., 1981).

Results and Discussion

Disease intensity

The results on the disease intensity of sheath blight due to the application of coir waste composted by *L. connatus*, *P. sajor - caju* and

urea, at different levels are presented in table 1. Among the mushroom fungi, *L. connatus* treated coir waste and urea at all levels significantly reduced the intensity of sheath blight, when compared with *P. sajor - caju*.

The treatments coir waste compost (*L. connatus*) @ 18.5 t + urea @ 60 kg N/ha and composted coir waste (*L. connatus*) @ 15.5 t/ha + urea @ 60 kg N / ha were found superior and significantly reduced the intensity of sheath blight (Gr. 3.00 and 3.52), compared to the existing fertilizer recommendation of 12.5t composted + 120 kg N/ha, where maximum disease intensity of Gr. 9.75 (sheath blight).The other treatments were on par with existing recommendation.

The similar trend was also observed with *P. sajor - caju*. The above said best treatments also significantly reduced the disease intensity of sheath blight (Gr. 5.60 and 6.20). The earlier treatments not significantly reduced the disease intensity and on par with existing recommendation (Fig. 1).

It is well known that lignocellulosic crop residues are the most abundant naturally occurring polymers in the biosphere. Fungi are unique among microorganisms in that they secrete a variety of extracellular enzymes to degrade lignocellulosic materials.


Decomposition of lignocellulosic residues is rated as the most important event in the carbon cycle of the earth (Bennett and Faison, 1997). Efficient and controlled degradation of lignocellulosic materials by these fungi lead to a number of processes of great economic importance. *Pleurotus* spp. well known white rot basidiomycotina produce lignocellulolytic enzymes to degrade complex of lignocellulosic residues, make their use as an alternative method presently used to overcome and reduce environmental pollution.

Table.1 Efficacy of coir waste compost on the intensity of sheath blight of rice (var. ADT 36)

S. No.	Fertilizer doses (ha)	* <i>L. connatus</i>	* <i>P. sajor-caju</i>
		Sheath blight	Sheath blight*
1.	1 2.5 t+ 120 kg N (control)	7.75	8.50
2.	12.5 t + 90 kg N	7.00	7.81
3.	12.5t + 60kgN	6.52	7.50
4.	15.5t + 90kgN	6.48	7.95
5.	15.5t + 60kgN	3.52	6.20
6.	18.5t + 90kgN	6.12	7.92
7.	18.5t + 60kgN	3.00	5.60
CD (P = 0.05)		1.74	1.60

	Sheath blight
Species	0.03
Treatment	0.02
Species x treatments	0.04

* Disease intensity on 0-9 scale



Coir waste compost on rice disease management

The current inorganic fertilizer prices and low availability of organic manure like farm yard manure, poultry manure etc., and their bulky need, limit the use of these available sources

of nutrients in Agriculture. Use of coir waste, which is available in huge quantities, composted with mushroom fungi have been reported to improve the nutritional status of the soil and reduced various plant diseases (Rajan, 1980; Theradimani and Marimuthu 1992; Sheela, 1992).

Increased in rice yield and significant reduction of major disease of rice by the application of 15.5 or 18.5 t of coir compost (*L. connatus*) and urea at 60 kg N per ha, might be due to low inorganic nitrogen and high organic crop residue supplementation. Increased inorganic nitrogen appears to influence disease potential than inoculum potential and hence, high inorganic nitrogen level leads to the outbreak of fungal disease (Baker and Martinson, 1970). Conversely, organic residues at high level decrease disease intensity, even though the population of the pathogen is very high (Nash and Synder, 1962; Huber and Watson, 1974). Moreover incorporation of coir waste composted by *L. connatus* which contained high amount of minerals might also an important role in the reduction of diseases intensity and improvement of yield (Agrios, 1997)

Increased in rice yield and significant reduction of major disease of rice by the application of 15.5 or 18.5 t of coir compost (*L. connatus*) and urea at 60 kg N per ha, might be due to low inorganic nitrogen and high organic crop residue supplementation. Increased inorganic nitrogen, appears to influence disease potential than inoculum potential and hence, high inorganic nitrogen level leads to the outbreak of fungal disease (Baker and Martinson, 1970). Conversely, organic residues at high level decrease disease intensity, even though the population of the pathogen is very high (Nash and synder, 1962; Huber and Watson, 1974). Moreover incorporation of coir waste composted by *L. connatus* which contained high amount of minerals, might also an important role in the reduction of diseases intensity and improvement of yield (Agrios, 1997).

References

Agrios, N., 1997. Environmental effects on the development of infectious plant

disease. In: Plant Pathology. Academic Press, pp. 144-151.

Amin, K.S., B.D. Sharma, and C.R. Das 1974. Occurrence in India sheath rot of rice caused by *Acrocyndrium oryzae*. *Pl. Dis. Reprtr*, 58: 358-360.

Anderson, D.C., A.T. Ralston, and J. Axelrod 1971. Chemical treatment of rye straw; *in vitro* dry matter digestibility and compositional changes. *J Animal Sci.*, 37: 148-152.

Anonymous, 1981. Annual Report for 1980-81. Int. Rice. Instt, Philipines. 165-169.

Bannett, J.W., and B.D. Faison 1997. Use of fungi in Biodegradation. In: Environmental microbiology, AISM, press, Washington Pp. 310-313.

Beg, S., S.I. Zafan, and F.H. 1986. Shah Rice husk bio degradation by *Pleurotus* species to produce a ruminant feed. *Agric wastes*, 17: 15-21.

Huber, D.M., and R.D. 1974. Watson, Nitrogen form and Plant Disease, In: Journal Series of Purdue University Agricultural Experimental Station, West Lafayette, Indiana. Pp. 139-165.

Lakshmanan, P., 2002. A new mushroom fungus identified in Tamil Nadu, *Indian J. Pl. Diseases and Protection*, (online).

Manickam, T.S., K. Seetharaman, K. Mayalagu and V. Mariappan 1980. Wealth from waste composted coir waste. *Mushroom Res*, 1: 45-47.

Marimuthu, T., 1995. Prospects of oyster mushroom in Tamil Nadu. *J Eco. Biol*, 7: 27-34.

Nash, S.M., and W.C. Synder 1962. Quantitative estimation of plate counts of propagules of the bean root rot *Fusarium* in field soils. *Phytopathology*, 52: 567-572.

Piper, C.S., 1966. Soil and Plant analysis. Hanos Publishers, Bombay, P.368

Rajan, K.M., 1980. Soil amendments in plant disease control. *IRRN*, 5, 15

Sheeba, R.I, M.A. Nair and B.K.

- Jayachandran 1999. Organic recycling in coconut. *Indian coconut J*, 49:10-12
- Sheela, J., 1992. Biological control of Fusarium wilt of *Solanum melongena* L. caused by *Fusarium solani* (Mart.) Sacc. M.Sc. (Ag.) Thesis, Tamil Nadu Agrl. Univ., Coimbatore, India. P.141.
- Singh, M.P., and L.J. Griensven 2000. Biodegradation of lignocellulosic waste through cultivation of *Pleurotus sajor-caju*. Proc. 15th Int. Cong. Sci. Culti. Edible Fungi, Maastricht, Netherlands, Pp. 617-618.
- Theradimani, M., and T. Marimuthu 1992. Utilization of *Pleurotus* spp. For decomposing coir pith. *Mushroom Res.*, 1: 49-52.
- Zadrazil, F., 1975. Influence of carbon dioxide concentration on the mycelial growth of three *Pleurotus* species. *Eur J Appl Microbio*, 1: 327-335.
- Zadrazil, F., 1984. Microbial conversion of lignocellulosic into feed. In: straw and other fibrous by products as feed. (Eds. S. Sundstal and Owen). Development in Animal and Veterinary Sciences, 14: 276-292.

How to cite this article:

Sudha, A., M. Senthilkumar and Vijayakumar, M. 2017. Efficacy of Coir Waste Compost and Urea at Different Levels on the Incidence of Sheath Blight of Rice. *Int.J.Curr.Microbiol.App.Sci*. 6(9): 778-783. doi: <https://doi.org/10.20546/ijcmas.2017.609.096>