

Review Article

<https://doi.org/10.20546/ijcmas.2019.806.154>

## Probiotic Fruit and Vegetable Juices: Approach Towards a Healthy Gut

Aditya Chaudhary\*

Department of Veterinary and Agricultural Sciences, Jayoti Vidyapeeth Women's University,  
Jaipur, India

\*Corresponding author

### ABSTRACT

#### Keywords

Probiotic fruit,  
Vegetable juices,  
Healthy gut

#### Article Info

Accepted:  
12 May 2019  
Available Online:  
10 June 2019

The useful effects of foods with added live microbes on human health are being increasingly promoted by health professionals. The major probiotic beverage available in the markets today, are milk-based products; usually in the form of fermented milk. However, with an increase in the consumer vegetarianism and demand for cholesterol free probiotics, there is also a demand for the non-dairy based probiotic beverage. Considering the above mentioned facts, fruits and vegetable juices may be the potential substrates, where the probiotic bacteria will make their mark, both in the developing and the developed countries. Fruits and vegetables are easily perishable commodities due to their high water activity and nutritive values. Lactic acid fermentation increases the shelf life of fruits and vegetables and also enhances various beneficial properties, including nutritive value and flavours, and reduces toxicity. As a whole, fermented fruit and vegetable juices with probiotics not only serve as food supplements but also attribute towards health benefits. This review aims at describing various factors affecting the viability of probiotic bacteria throughout the storage period in diverse fruit and vegetable juices and perspective technologies to improve the survivability of probiotics.

### Introduction

Today, all consumers have a considerable amount of their nutritional needs met through fermented foods and beverages (Steinkraus, 1997). Fermented/ functional foods are those containing bioactive compounds viz. dietary fibers, oligosaccharides, minerals, vitamins and probiotics (active friendly bacteria), having pleasant aroma, flavour and texture, that maintain the equilibrium of intestinal microflora (Steinkraus, 1997; Jankovic *et al.*, 2010; Shah and Prajapati, 2007). Fermented foods and beverages have diversity in cultural

preferences and traditions found in the different geographical areas, where they are produced. Fermentation has helped the ancestors in colder regions to survive during the winter season and those in the tropics to survive drought periods (Swain *et al.*, 2014). According to FAO (1998), fermentation is a slow decomposition process of organic substances where carbohydrates are converted to alcohols or organic acids with the help of microorganisms or their enzymes. Fermentation processes have been developed for preserving the fruits and vegetables by preserving them by organic acids or alcohols,

impart the desired flavour and texture to the food, decrease the cooking time and also reduce the toxicity (Rolle, 2002). Lactic acid fermentation of fruits and vegetables is a usual practice for maintaining and improving the nutritional and sensory attribute of the food products (Rolle and Satin, 2002; Endrizzi *et al.*, 2009; Demir *et al.*, 2006). Certain nutrients such as minerals and vitamins and acidic nature of fruits and vegetables provide a favourable medium for fermentation by Lactic acid bacteria. Probiotics are defined as live microbial feed, which is supplemented by food that maintains the intestinal health of the host in a beneficial manner (Tamang, 2009). Various studies suggest that supplementation of probiotic via food has several health benefits such as enhanced immune system (McNaught and MacFie, 2001), improved gastrointestinal function (Saarela *et al.*, 2002), reduction of serum cholesterol (Anandharaj *et al.*, 2014) and lower the risk of colon cancer (Rafter, 2003).

World Health Organization (WHO) and Food and Agriculture Organization (FAO) approved a specific daily intake dose of fruits and vegetables to prevent chronic pathologies such as coronary heart disease, hypertension, and risk of strokes (Swain *et al.*, 2014). Nowadays, the consumer prefers to intake the food and beverages, which is fresh, highly nutritional, ready to eat or drink and health-promoting (Endrizzi *et al.*, 2009). According to many literatures, the functional food market constitutes one of the main captivating areas of investigation and innovation in the food sector. As suggested by one survey, probiotics will be incorporated in various food products viz. dietary supplements, functional foods, animal feed, and in therapeutic and preventive healthcare applications and the probiotic market will rise to worth \$46.55 Billion by 2020 (Anon, 2016). Plant based foods in Asian diets is the

main food of the daily intake as they consume a relatively low amount of meat and dairy foods. Excluding dietary habits, lactose intolerance prevents many Asian people from consuming milk. In last decade, the average per capita dairy consumption in milk equivalent for the major dairy markets was 97.6 Kg in Japan, 80 Kg in South Korea, 71.8 Kg in India, 67.8 Kg in Malaysia, 28.7 Kg in Thailand, 24 Kg in Philippines, 10.2 Kg in China, 8.6 Kg in Vietnam and 7.8 Kg in Indonesia. This creates disparity with per capita consumption of 330 Kg in EU-15, 310 Kg per capita in Australia, and 251 Kg per capita in the United States (Dong, 2006). Therefore, in view of the above facts, fruits and vegetables can be utilized as the potential substrates, where healthy probiotic bacteria will make their impression, both in the developing and the developed countries.

### **Probiotics in fruit and vegetable juices**

Hippocrates once wrote: 'Let food be thy medicine, and let medicine be thy food.' Nowadays, food having medicinal value has been renewed as 'functional foods.' A probiotic may also be a functional food (Scheinbach, 1998). Functional foods are defined as 'foods that contain traditional nutrition along with some health-promoting component(s).' Functional foods are also known as designer foods, nutraceuticals, therapeutic foods, medicinal foods, foodiceuticals, medifoods, and superfoods (Soccol *et al.*, 2010). In general, the term functional foods mentions to a food that has been improved in some way to become 'functional.' One way in which foods can be improved to become functional is by the incorporation of probiotics (FAO/WHO). Food products have been formulated with the addition of probiotic cultures. Different food matrices have been used such as various types of cheeses, powdered milk, ice-creams, milk-based desserts, butter, mayonnaise, and

fermented food of vegetable origin (Tamime *et al.*, 2005). Presence of allergens, lactose intolerance, fat content, and requirement of cold storage are some limitations of probiotic dairy products (Heenan *et al.*, 2004; Yoon *et al.*, 2006). This aspect has led to the initiation of novel probiotic products based on non-dairy matrices (Table 1). In recent times, various raw materials have been investigated extensively for determining the appropriate matrix to produce new non-dairy functional foods (Vasudha and Mishra, 2013). Matrices used in the development of non-dairy probiotic products are fruits, vegetables, cereals, and legumes. Fruits and vegetables are considered good matrices as they contain nutrients such as minerals, dietary fibers, vitamins, and antioxidants (Patel, 2017) while lacking the dairy allergens that might prevent consumption by a particular section of the population (Luckow and Delahunty, 2004). The development of various fruit juices has been studied (Yoon *et al.*, 2004; Soccol *et al.*, 2007).

Besides potential sensory challenges, there is a sincere attentiveness towards the development of fruit and vegetable-based functional juice/ beverage, fortified with probiotic and prebiotic ingredients. The fruit and vegetable-based juices inherently contain beneficial nutrients, they have taste profiles that are appealing to all age groups, they are recognized as being refreshing and healthy, and therefore, they have been recommended as a suitable medium for the functional health ingredients (Tuorila and Cardello, 2002).

Hardaliye is a lactic acid fermented beverage, prepared by the natural fermentation of red grape juice supplemented with crushed mustard seeds and benzoic acid. This drink is originated in the Thrace region of Turkey. Eteric oils of mustard seeds affect the yeast and give a characteristic flavour to the final product. Benzoic acid also affects the yeast by

inhibiting or decreasing the production of alcohol. Hardaliye is stored at 4°C after fermentation. It can be consumed fresh or aged. The lactic acid bacteria found in this beverage are *Lactobacillus acetotolerans*, *Lactobacillus brevis*, *Lactobacillus casei* subsp. *pseudoplantarum*, *Lactobacillus paracasei* subsp. *paracasei* and *Lactobacillus vaccinoferus* (Arici and Coskun, 2001). Globally, there are so many commercially available probiotic fruit or vegetable juices, mentioned in Table 2.

### **Fermentation of fruit and vegetable juice by probiotic bacteria**

The consumption of LA fermented vegetable juices (lacto-juices) has increased in many countries. Lacto-juices are produced mainly from cabbage, carrot, celery, red beet, and tomato (Demir *et al.*, 2006). They can be produced by either of the following ways:

- (i) Fermentation of vegetable mash or juice;
- (ii) Usual way of vegetable fermentation and then processed by pressing the juice.

There are three ways of lactic fermentation of fruit and vegetable juices:

- (i) Spontaneous fermentation by natural microflora;
- (ii) Fermentation by starter cultures (added to the raw materials);
- (iii) Fermentation by heat-treated materials by starter cultures.

During the production of lacto-juices, the pressed juice can be sterilized at first and consecutively it is inoculated by a starter culture mainly a LAB at a concentration varying from  $2 \times 10^5$  to  $5 \times 10^6$  CFU/ml (Demir *et al.*, 2006; Panda and Ray, 2007). The difference between the production of

fermented and non-fermented juice is explained in Figure 1. For fermentation of juices of high quality, it is necessary to use commercially supplied starter cultures viz. *L. plantarum*, *L. bifidus*, *L. brevis*, *L. xylosus*, and *L. bavaricus*. The criteria used for determining the suitability of a strain are mentioned below (McFeeters, 2004):

- (i) Ability of a substrate to acquire a starter culture;
- (ii) The rate and total production of lactic acid, change in pH, loss of nutritionally important substances;
- (iii) Decrease in nitrate concentration and production of biogenic amines;
- (iv) Type of metabolism and ability of culture to produce desirable sensory properties of fermented products.

### **Major factors affecting the survivability of probiotic bacteria in juices**

The health benefits of probiotic juices mainly depend on the concentration of probiotics in the juices, along with their ability to survive the unfavourable conditions of the gastrointestinal tract. Two important criteria to be fulfilled by fruit juices are to maintain the viability (at least  $10^6 - 10^7$  cells/ml) and activity of probiotics in the final product at the end of the shelf life (Nualkaekul and Charalampopoulos, 2011).

However, probiotic viability is strain dependent (Corbo *et al.*, 2012; Tripathi and Giri, 2014; Do Espírito Santo, 2011), i.e., some strains of *L. plantarum*, *L. acidophilus* and *L. casei* can grow in fruit matrices as they survive in acidic environments (Peres *et al.*, 2012). Although, fruit and vegetable juices have presented limitations to the addition of probiotic cultures such as (1) Intrinsic food parameters – pH, titratable acidity, molecular oxygen, water activity, presence of sugar, salt, artificial flavouring and colouring agent and

chemical and microbial preservatives like bacteriocins and hydrogen peroxide; (2) processing parameters – incubation temperature, extent of heat treatment, cooling rate; (3) microbiological factors – kind of probiotic strains, rate and proportion of inoculum, inter-compatibility of different strains.

Probiotic cultures are neutrophils, with a pH optimum between 5 and 9. When the cells are present in low pH environments, there is an increased requirement of energy for the maintenance of intracellular pH and ATP is lacking for other crucial functions, resulting in cell death (Nualkaekul *et al.*, 2011). Also, they are usually microaerophilic or anaerobic and therefore, they lack electron carrier chain and the catalase enzyme, and the presence of oxygen may cause formation and accumulation of toxic metabolites in the cell and cause cell death from oxidative damage. Furthermore, the presence of preservatives, flavourings, and colourings may decrease the survivability of probiotic cultures. The probiotic cultures can change the sensory attributes (development of unpleasant aroma and flavour).

The metabolism of probiotic lactic acid bacteria may result in the production of metabolites and can affect the flavour and aroma of the food product. Among various probiotics, lactobacilli were found to survive in fruit juices with pH ranging 4.3 to 3.7; whereas bifidobacteria are less acid-tolerant (Tripathi and Giri, 2014).

Therefore, the type of microorganism and juice, storage conditions, and the addition of other compounds may influence the sensory traits of the finished product. Fermented juices with sugars had a more accepted taste and flavour than the sugar-free juices (Sivudu *et al.*, 2014).

## **Improving probiotic survival in juices**

### **Fortification with prebiotics**

The simplest way to improve the viability of probiotics in fruit juices could be the fortification with some prebiotics (dietary fibers, cellulose) or with some elements able to exert a protective effect within the juice. Apple juice incorporated with *L. rhamnosus* was fortified with oat flour, and 20% of  $\beta$ -glucan, and this could protect the probiotic bacteria during refrigerated storage (Saarela *et al.*, 2006). Orange juice fortified with pearl millet exhibited high viability of *L. plantarum* and a good source of fibers that contribute to maintaining the intestinal balance and had the potential to exhibit prebiotic effects such as the effects on the sensory traits and post acidification (Gupta and Sharma, 2016). Beetroot and carrot juices were enriched with brewer's yeast autolysate before lactic acid fermentation with *L. acidophilus*. The supplementation with autolysate convincingly increased the number of lactic acid bacteria during fermentation (Aeschlimann and Von Stocar, 1990), decreased the fermentation time (Chae *et al.*, 2001). Use of brewer's yeast helped in the economic optimization of the fermentation. A mixture of beetroot and carrot juice had an optimum quantity of antioxidants, minerals, pigments, amino acids, and vitamins after the addition of brewer's yeast (Rakin *et al.*, 2007). Fig juice incorporated with *L. delbrueckii* and fortified with inulin had a high viable count than the control (probiotic fig juice) and exhibited high antioxidants, polyphenols, medicinal properties, and organoleptic attributes (Khezri *et al.*, 2018)

### **Induction and adaptation of resistance**

Exposure of probiotics to sub-lethal stress could instigate a resistance and an adaptive stress response (Gobetti *et al.*, 2010). Many

researchers examined this approach. Viability of *L. reuteri* DSM 20016 was examined in pineapple, orange, green apple, and red fruit juices and found that the viability of the probiotic strain lost due to the combined effect of pH and phenols. Two different strategies were executed to overcome this survivability loss. Strain cultivation in different concentrations of red fruit juices (upto 50%) and provide phenol stress in the form of addition of vanillic acid and acid stress by acidification to pH 5.0. These approaches resulted in the prolongation of the viability of *L. reuteri* by 5 days (phenol stress) and 11 days (acid stress) (Perricone *et al.*, 2014). Survivability of *B. breve* in a blended juice of orange-grape and passion fruit was improved by creating an acid tolerance variant of the bacteria by UV mutagenesis along with cultivation at sub-lethal pHs (Saarela *et al.*, 2011).

### **Storage under refrigeration and microencapsulation**

Lactic acid bacteria are highly sensitive to variation in storage temperature. The probiotic strain viability in fruit or vegetable juice is found to be affected, as refrigeration could promise a more prolonged survival, whereas thermal abuse could exhibit a detrimental effect (Patel, 2017). Many authors proposed various approaches to resolve the issue of thermal abuse. Microencapsulation techniques have been successfully applied using different matrices to protect the probiotic bacterial cell from the damage caused by the external environmental components (Patel, 2017). A novel microencapsulation method helped in reducing the acidification and improving the viability of probiotic strains *L. acidophilus* and *L. rhamnosus* at 25°C for 9 days in orange juice (Sohail *et al.*, 2012). The cells of *B. animalis* subsp. *lactis* Bb12 were microencapsulated in a milk protein matrix.

The stability of free and encapsulated probiotic cells was studied in pineapple juice and strawberry-apple juice during 28-day storage. Pineapple juice was proved to be a suitable carrier for encapsulated probiotic cell viability whereas microcapsules and free cells were not stable in strawberry-apple juice and died after 14 days of storage at 22°C (Horáčková *et al.*, 2018). *L. acidophilus* immobilized in Ca-alginate exhibited a higher cell survival rate in tomato juice and overall acceptance by the sensory evaluation when compared with the free cells during the cold storage at 4°C (King *et al.*, 2007). A novel probiotic food product had been prepared with *L. acidophilus* when immobilized with Ca-alginate and found to carry out the normal banana puree fermentation (Tsen *et al.*, 2004). Thai herbal extracts such as cashew flower, pennywort, and yanang were alginate encapsulated and studied for the viability of probiotic bacterial isolates *L. casei* 01b, *L. acidophilus* LA5 and *B. lactis* Bb-12 in mulberry, maoberry, longan, and melon juices. After 30 days of cold storage, higher viability of *L. casei* 01b entrapped with 0.05% (w/v) cashew flower extract was determined as compared with those encapsulated with pennywort and yanang extracts (Chaikham, 2015). Microencapsulated probiotic cells in fruit juices were found to be more stable in comparison with free probiotic cells. The encapsulated probiotic cell of *L. acidophilus*, *L. paracasei*, *L. plantarum*, *L. rhamnosus*, *L. salivarius*, *B. longum*, *B. lactis* Bi-04, and Bi-07 managed to survive in the acidic environment of the orange juice, did not exhibit a survivability loss, and 5 log CFU/ml cell count was determined (Ding and Shah, 2008). Viable count of encapsulated *L. plantarum* in grapes, orange, sapodilla, and watermelon juices was evaluated as 7 log CFU/ml or more (Gaanappriya *et al.*, 2013). It was reported that microencapsulation provides a favourable anaerobic environment for sensitive probiotic bacteria and provide a

physical hindrance from the harsh acidic conditions of the fruit or vegetable juices (Ding and Shah, 2008). The probiotic cultures of *L. casei*, *L. plantarum*, *L. fermentum*, *Lysinibacillus sphaericus*, and *Saccharomyces boulardii* were microencapsulated using alginate coated chitosan beads and incorporated in tomato and carrot juices. The juices were pasteurized for 20 min at 63°C and incubated at 37°C for 72 h. Viable cell count of *Lysinibacillus sphaericus* and *Saccharomyces boulardii* were increased from 6.5 to 8.9 log CFU/ml and 5.2 to 7.6 log CFU/ml, respectively during 24-42 h and found to be decreased afterward (Naga *et al.*, 2016).

### **Use of antioxidants**

During the storage of packaged foods, the level of oxygen should be maintained low in order to avoid oxidative damage to the probiotics, however, the degree of sensitivity is highly strain variable. Oxidative damage by oxygen is due to the generation of ROS (Reactive Oxygen Species) such as hydrogen peroxide or superoxide ion. Bifidobacteria are more sensitive than lactic acid bacteria (Nag and Das, 2013). Modification of the product atmosphere was suggested by many authors by raising the CO<sub>2</sub> content in the headspace (Corbo *et al.*, 2014). Incorporation of antioxidants could help to control the harmful effects of oxygen. The probiotic bacteria HOWARU *L. rhamnosus* HN001, HOWARU *B. lactis* HN001 and *L. paracasei* LPC 37 were incorporated into model juice along with vitamins viz. vitamin B2, vitamin B3, vitamin B6, vitamin C and vitamin E and antioxidants such as grape seed and green tea extract. The model juice containing grape seed extract, green tea extract, and vitamin C showed better survival of probiotic bacteria, exhibited viability of 4.29 log CFU/ml, 7.41 log CFU/ml, and 6.44 log CFU/ml after 6 weeks of storage (Shah *et al.*, 2010).

Effects of different amounts of (+)-catechin, green tea extracts, and green tea epigallocatechin gallate were studied on the viability of *B. longum* ATCC 15708, *B. longum* subsp. *infantis* ATCC 15697 and *L.*

*helveticus* R0052 having different oxygen sensitivities and found that the growth of *L. helveticus* was greatly enhanced (Gaudreau *et al.*, 2013).

**Table.1** The diversity of probiotic food products and the viability of each probiotic in different products at the end of appropriate storage conditions


Product type (Fruit and vegetable based)	Probiotic strain	Viability at the end of storage	Total storage time	References
Apple juice	<i>L.acidophilus</i> <i>L.rhamnosus</i> <i>L.plantarum</i> <i>L.salivarius</i> <i>L.paracasei</i> <i>B.longum</i> <i>B.lactis</i> type Bi-04 <i>B.lactis</i> type Bi-07	10 <sup>6</sup> CFU/mL	6 weeks	Ding and Shah, 2008
Amla ( <i>Phyllanthusemblica</i> )	<i>L.paracasei</i> HII01			Peeranjan <i>et al.</i> , 2016
Beet root juice	<i>L.plantarum</i> C3, <i>L.casei</i> A4, <i>L.acidophilus</i> LA39, <i>L.delbrueckii</i> D7	10 <sup>6</sup> – 10 <sup>8</sup> CFU/mL	4 weeks	Yoon <i>et al.</i> , 2005
Cabbage juice	<i>L.plantarum</i> C3, <i>L.casei</i> A4 <i>L.delbrueckii</i> D7	4.1 X10 <sup>7</sup> CFU/mL ( <i>L.plantarum</i> ), 4.5 X 10 <sup>5</sup> CFU/mL ( <i>L.delbrueckii</i> ), <i>L.casei</i> lost complete cell viability after 2 weeks of	4 weeks	Yoon <i>et al.</i> , 2006
Carrot juice	<i>B.lactis</i> Bb-12, <i>B.bifidum</i> B7.1 & B3.2	15-17 mg/ml lactic acid production	4 weeks	Kun <i>et al.</i> , 2008
Carrot juice	<i>L.rhamnosus</i> , <i>L.bulgaricus</i> with inulin or fructooligosaccharides		4 weeks	Nazzaro <i>et al.</i> , 2008
Carrot, beet and apple juice	<i>L.casei</i>	1.5 x 10 <sup>6</sup> CFU/ml	4 weeks	Zandiet <i>al.</i> , 2016
Carrot blended with	<i>L.plantarum</i> CECT	10 <sup>8</sup> – 10 <sup>9</sup>	30 days	Valero-Cases and

orange juice	220	CFU/MI		Frutos, 2017
Fermented Kale juice	<i>L.plantarum</i> , <i>L.casei</i> , <i>L.acidophilus</i> , <i>L.brevis</i>	10 <sup>8</sup> CFU/ml	4 weeks	Kim, 2017
Fermented vegetable soybean beverage	<i>L.acidophilus</i> La-5, <i>B.animalis</i> Bb-12	~10 <sup>6</sup> CFU/MI 10 <sup>8</sup> CFU/MI	28 days	Battistini <i>et al.</i> , 2017
Mango, sapota, grape and cantaloupe	<i>L.casei</i>		4 weeks	Kumar <i>et al.</i> , 2013
Noni juice	<i>L.casei</i> , <i>L.plantarum</i> , <i>B.longum</i>	10 <sup>8</sup> CFU/MI	4 weeks	Wang <i>et al.</i> , 2009
Orange juice after spouted bed drying	<i>L.casei</i>	~10 <sup>6</sup> CFU/g	5 weeks	Alves, 2017
Orange, pineapple and cranberry juice	<i>L.casei</i> , <i>L.paracasei</i> , and <i>L.rhamnosus</i>	10 <sup>8</sup> CFU/ml (Orange Juice), 10 <sup>7</sup> CFU/ml (Pineapple juice) and very low cell viability was seen in cranberry juice	12 weeks	Sheehan <i>et al.</i> , 2007
Peach Juice	<i>L.casei</i> , <i>L.delbrueckii</i>	1.72 x 10 <sup>7</sup> CFU/ml	4 weeks	Pakbinet <i>et al.</i> , 2014
Pineapple juice	<i>L.casei</i>	10 <sup>6</sup> CFU/ml	42 days	Costa <i>et al.</i> , 2013
Pomegranate juice	<i>L.plantarum</i> <i>L.paracasei</i> <i>L.acidophilus</i> <i>L.delbrueckii</i>		4 weeks	Mousaviet <i>et al.</i> , 2011
Red grape and white grape juice	<i>P.pentosaceus</i>	3.2 – 6.5 log CFU/ml	72 h	Kumar <i>et al.</i> , 2017
Sugarcane and Sweet lime juices	<i>L.acidophilus</i>	2.0 x 10 <sup>8</sup> – 5.5 x 10 <sup>8</sup> CFU/ml	3 weeks	Khatoon and Gupta, 2015
Watermelon and tomato juice	<i>L.fermentum</i> , <i>L.casei</i>		4 weeks	Sivudu <i>et al.</i> , 2014
Tomato juice	<i>L.plantarum</i> C3, <i>L.casei</i> A4, <i>L.acidophilus</i> LA39, <i>L.delbrueckii</i> D7	10 <sup>6</sup> – 10 <sup>8</sup> CFU/mL	4 weeks	Yoon <i>et al.</i> , 2004
Tomato, orange and grape juices	<i>L.plantarum</i> , <i>L.acidophilus</i>			Nagpal <i>et al.</i> , 2012

**Table.2** Commercially available probiotic juices (Adapted from Patel, 2017; Molin, 2001; Leporanta, 2005)

Product name	Manufacturer	Probiotic strain(s)	Major attributes
<b>BiolaR</b>	TINE BA, Norway	<i>L.rhamnosus</i> GG	Mixture of apple-pear and, orange-mango juice with 95% fruit and no added sugar
<b>Bio-Live Gold &amp; Dark</b>	Bio-Live/Microbz Ltd., UK	Mixtures of 13 strains including <i>L.acidophilus</i> , <i>L.bulgaricus</i> , <i>L.casei</i> , <i>L.plantarum</i> , <i>L.fermentum</i> , <i>Lactococcus lactis</i> , <i>Bacillus subtilis</i> , <i>B.bifidum</i> , <i>B.longum</i> , <i>B.infantis</i> , <i>Streptococcus thermophilus</i> , <i>Comobcillus</i> , and <i>Saccharomyces cerevisiae</i>	Mixture of fruit juices such as acai berry, cherry, goji, noni, pomegranate, lemon, and various herbs
<b>Bravo Friscus</b>	Probi AB, Sweden	<i>L.plantarum</i> HEAL9 and <i>L.paracasei</i> 8700:2	Orange apple and tropical fruit juices
<b>Gefilus Fruit drinks</b>	Valio Ltd., Finland	<i>L.rhamnosus</i> GG and <i>Propionibacterium freudenreichii</i> ssp. <i>shermanii</i> JS	Berries
<b>Golden Circle Healthy Life Probiotic Juice</b>	Golden Circle, Australia	<i>L.paracasei</i> 8700:2 and <i>L.plantarum</i> HEAL9	Mixture of apple juice with mango puree or orange, apple, pineapple, passion fruit with banana puree
<b>Goodbelly® Carrot Ginger Flavor</b>	Goodbelly, USA	<i>L.plantarum</i> 299v (50 million cells/ portion)	Carrot juice, ginger extract, and cane sugar contains 2% or less gluten-free oat flour
<b>KEVITA</b>	KEVITA, USA	<i>L.rhamnosus</i> , <i>L.plantarum</i> , <i>L.paracasei</i> , <i>Bacillus coagulans</i> GBI-30 6086	Various fruit based mixtures such as strawberry and coconut, lime, mint and coconut, pineapple and coconut
<b>Malee Probiotics</b>	Malee Enterprise Company Ltd., Thailand	<i>L.paracasei</i>	Prune, grape and orange juice
<b>PERKii Probiotic Water</b>	PERKii, Australia	<i>L.paracasei</i> Lc431	Fruit juice mixtures such as raspberry and pomegranate, lime and coconut, mango and passion fruit and strawberry and watermelon
<b>Probiotic Naked Juice</b>	Naked® Juice, USA	<i>Bifidobacterium</i>	Mixture of apple, orange, pineapple juices and mango and banana puree with fructooligosaccharides
<b>ProViva</b>	EMEA Probi AB, Sweden	<i>L.plantarum</i> 299v (50 million cells/ portion)	Orange, strawberry or blackcurrant juice and fortified with 5% oat flour
<b>Rela Fruit Juice</b>	Biogaia Global, Sweden	<i>L.reuteri</i> MM53	Fruit juice
<b>Tropicana probiotics</b>	Tropicana, USA	<i>B.lactis</i>	Fruit juice mixtures such as strawberry and banana, pineapple and mango and peach passion fruit
<b>Vita Biosa</b>	Biosa, Denmark	<i>B.lactis</i> , <i>B.longum</i> , <i>L.acidophilus</i> , <i>L.casei</i> , <i>L.rhamnosus</i> , <i>L.salivarius</i> , <i>L.lactis</i> , <i>Streptococcus thermophilus</i>	Ginger, blackcurrant, blueberry juice with ginger extract enriched with 19 different aromatic herbs

**Fig.1** Production of fermented and non-fermented fruit/vegetable juice enriched with probiotics



In conclusion, most probiotic products available in the market are milk-based; however, the increase in the consumer vegetarianism and demand for cholesterol free probiotics has led researchers and scientists to explore newer matrices as vehicles for probiotics. Technological advances have made it possible to alter some structural characteristics of fruit and vegetable matrices by modifying food components in a controlled way. Fruits and vegetables make the ideal substrates for the culture of probiotics since they already contain beneficial nutrients such as antioxidants, minerals, vitamins, and dietary fibers while

lacking the dairy allergens that might prevent the consumption by certain groups of the population. Vegetables and fruits reported containing a wide variety of antioxidant components, including phytochemicals.

There is a genuine interest in the development of fruits and vegetable functional beverage with probiotics because they serve as a healthy alternative for dairy probiotics, are cholesterol free and also favour consumption by lactose tolerant consumers. Detailed studies on the microbial composition and characteristics of fermented fruits and vegetable juices lead to further application.

## References

- Aeschlimann, A. and Von Stocar, U. 1990. The effect of yeast extract supplementation on the production of lactic acid from whey permeate by *Lactobacillus helveticus*. *Applied Microbiology and Biotechnology*. 32, 398–440.
- Alves, N.N. 2017. Spouted bed as an efficient processing for probiotic orange juice drying. *Food Res. Int.* 101, 54–60.
- Anandharaj, M., B. Sivasankari, and R. P. Rani. 2014. Effects of probiotics, prebiotics, and synbiotics on hypercholesterolemia: a review. *Chinese Journal of Biology*, vol. 2014, Article ID 572754, 7 pages, 2014.
- Anon. (2016) Retrieved on June 11, 2016 from FSIS. Website: <http://www.marketsandmarkets.com/PressReleases/probiotics.asp>.
- Arici, M. and Coskun, F. 2001. Hardaliye: Fermented grape juice as a traditional Turkish beverage. *Food Microbiology*. 18, 417–421.
- Battistini, C., Gullon, B., Ichimura, E.S., Gomes, A.M.P., Ribeiro, E.P., Kunigk, L., Moreira, J.U.V., Jurkiewicz, C.J. 2017. Development and characterization of an innovative synbiotic fermented beverage based on vegetable soybean. *Braz. J. Microbiol.* in press.
- Carlos Ricardo Soccol, Luciana Porto de Souza Vandenberghe, Michele RigonSpier, Adriane Bianchi Pedroni Medeiros, Caroline Tiemi Yamaguishi, Juliano De Dea Lindner, Ashok Pandey and Vanete Thomaz-Soccol. 2010. The potential of Probiotics. *Food Technol. Biotechnol.* 48(4): 413-434.
- Chae, H.J., Joo, H. and In, M. J. 2001. Utilization of brewer's yeast cells for the production of food grade yeast extract. Part I: Effects of different enzymatic treatments on solid and protein recovery and flavor characteristics. *Bioresource Technology*. 76: 253–258.
- Chaikham, P., 2015. Stability of probiotics encapsulated with Thai herbal extracts in fruit juices and yoghurt during refrigerated storage. *Food Bioscience*. 12: 61– 66.
- Corbo MR., Bevilacqua A, Petruzzi L, Casanova FP, Sinigaglia M. 2014. Functional beverages: The emerging side of functional foods commercial trends, research, and health implications. *Compr. Rev. Food Sci. Food Saf.* 13, 1192-1206.
- Costa, M.G.M., Fonteles, T.V.; de Jesus, A.L.T.; Rodrigues, S. 2013. Sonicated pineapple juice as substrate for *L. casei* cultivation for probiotic beverage development: Process optimisation and product stability. *Food Chem.*, 139, 261–266.
- de Vrese M., Stegelmann A, Richter B, Fenselau S, Laue C, Schrezenmeir J. 2001. Probiotics-compensation for lactase insufficiency. *American Journal of Clinical Nutrition*. 73(2): 421S–429S.
- Demir, N., K. S. Bahçeci, and J. Acar. 2006. The effects of different initial *Lactobacillus plantarum* concentrations on some properties of fermented carrot juice. *Journal of Food Processing and Preservation*. 30(3): 352–363.
- Demir, N., K. S. Bahçeci, and J. Acar. 2006. The effects of different initial *Lactobacillus plantarum* concentrations on some properties of fermented carrot juice. *Journal of Food Processing and Preservation*. 30(3): 352–363.
- Ding, W.K., and Shah, N.P. 2008. Survival of free and microencapsulated probiotic bacteria in orange and apple juices. *Int. Food Res. J.* 15, 219–232.
- Do Espírito Santo, A.P., Perego, P., Converti, A., Oliveira, M.N. 2011. Influence of food matrices on probiotic viability-A review focusing on the fruity bases. *Trends Food Sci. Technol.* 22, 377–385.
- Endrizzi I., G. Pirretti, D.G. Cal'ò, and F. Gasperi. 2009. A consumer study of fresh juices containing berry fruits. *Journal of the Science of Food and Agriculture*. 89(7): 1227–1235.
- FAO, Fermented Fruits and Vegetables-A

- Global Perspective, FAO Agricultural Services Bulletin, Rome, Italy, 1998, 134.
- Food and Agriculture Organization/World Health Organization (FAO/WHO), Guidelines for the evaluation of probiotics in food, Report of a Joint FAO/WHO Working Group on Drafting Guidelines for the Evaluation of Probiotics in Food, London, Ontario, Canada (2002) (<http://ftp.fao.org/es/esn/food/wgreport2.pdf>).
- Gaanappriya, M., Guhankumar, P., Kiruththica, V., Santhiya, N. and Anita, S. 2013. Probiotication of fruit juices *Lactobacillus acidophilus*. *International Journal of Advanced Biotechnology Research*. 4, 72–77.
- Gaudreau, H., Champagne, C.P., Remondetto, G.E., Bazinet, L. and Subirade, M. 2013. Effect of catechins on the growth of oxygen-sensitive probiotic bacteria. *Food Research International*. 53, 751–757.
- Gobbetti, M., di Cagno, R., de Angelis, M. 2010. Functional microorganisms for functional food quality. *Crit. Rev. Food Sci. Nutr*. 50, 716–727.
- Heenan, C.N., Adams, M. C., Hosken, R. W. and Fleet, G. H. 2004. Survival and sensory acceptability of probiotic microorganisms in a nonfermented frozen vegetarian dessert. *Lebensmittel-Wissenschaft und-Technology*. 37, 461–466.
- J. Karovicova and Z. Kohajdov´a. 2003. Lactic acid fermented vegetable juices,” *Horticultural Science*. 30, 152–158.
- J. Rafter. 2003. Probiotics and colon cancer,” *Bailliere’s Best Practice and Research in Clinical Gastroenterology*. 17(5): 849–859.
- Jankovic, I., Sybesma, W., Phothirath, P., Ananta, E. and Mercenier, A. 2010. Application of probiotics in food products-challenges and new approaches. *Current Opinions in Biotechnology*. 21, 175–181.
- Jayawardana, N., W.I.A.; Prasanna, P.H.P.; Ranadheera, C.S.; de Zoysa, H.K.S.; Vidanarachchi, J.K. 2015. Probiotics in functional foods. In *Fermented Foods: Sources, Consumption and Health Benefits*; Nova Science Publishers: New York, NY, USA, 103–126.
- K.Y. Yoon, E.E. Woodams, Y.D. Hang. 2004. Probiotication of tomato juice by lactic acid bacteria. *J. Microbiol*. 42, 315–318.
- Khatoon, N., and Gupta, R.K. 2015. Probiotics beverages of sweet lime and sugarcane juices and its physiochemical, microbiological and shelf life studies. *Journal of Pharmacognosy and Phytochemistry*. 4(3): 25-34
- Kim, S.Y., 2017. Production of Fermented Kale Juices with *Lactobacillus* Strains and Nutritional Composition. *Prev. Nutr. Food Sci*. 22, 231–236.
- King, V.A.E., Huang, H.Y. and Tsen, J.H. 2007. Fermentation of tomato juice by cell immobilized *Lactobacillus acidophilus*. *Mid-Taiwan Journal of Medicine*. 12, 1–7.
- Kumar, B.V., Mannepula S., Obulam V. and Reddy S. 2013. Physico-chemical analysis of fresh and probioticated fruit juices with *Lactobacillus casei*. *International Journal of Applied Science and Biotechnology*. 1(3):127-131.
- Kumar, B.V., Vijayendra, S.V.N.; Reddy, O.V.S. 2015. Trends in dairy and non-dairy probiotic products—A review. *J. Food Sci. Technol*. 52, 6112–6124.
- Kun, S., Rezessy Szabo, J.M., Nguyen, Q.D. and Hoschke, A. 2008. Changes of microbial population and some components in carrot juice during fermentation with selected *Bifidobacterium* strains. *Process Biochemistry*. 43(8): 816-821.
- Levri KM, Ketvertis K, Deramo M, Merenstein JH, D’Amico F. 2005. Do probiotics reduce adult lactose intolerance? A systematic review. *Journal of Family Practice*. 54(7):613–620.
- Luckow, T., and Delahunty, C. 2004. Which juice is ‘healthier’? A consumer study of probiotic non-dairy juice drinks. *Food Quality and Preference*. 15, 751–759.
- Manas Ranjan Swain, Marimuthu Anandharaj,

- Ramesh Chandra Ray, and Rizwana Parveen Rani. 2014. Fermented Fruits and Vegetables of Asia: A Potential Source of Probiotics. *Biotechnology Research International*, Article ID 250424, 19 pages.
- McNaught, C.E., and J. MacFie. 2001. Probiotics in clinical practice: a critical review of the evidence. *Nutrition Research*. 21(1): 343–353.
- Mohammad Mehdi Zandi, Mahnaz Hashemiravan and Shila Berenji. 2016. Production of probiotic fermented mixture of carrot, beet and apple juices. *Journal of Paramedical Sciences*. 7(3): 17-23.
- Molin, G. 2001. Probiotics in foods not containing milk or milk constituents, with special reference to *Lactobacillus plantarum* 299v. *American Journal of Clinical Nutrition*. 73, 380–385.
- Mousavi Z.E., Mousavi S.M. Razavi S.H., Emam-Djomeh Z. and Kiani H. 2011. Fermentation of pomegranate juice by probiotic lactic acid bacteria. *World Journal of Microbiology and Biotechnology*. 27(1):123-128.
- Mridula Gupta and Somesh Sharma. 2016. Prebiotics fortified with fruit juices- A good carrier for probiotics. *International Journal of Bio-Technology and Research*. 6(1): 43-48.
- Nag, A., and Das, S. 2013. Improving ambient temperature stability of probiotics with stress adaptation and fluidized bed drying. *Journal of functional foods*. 5,170–177.
- Naga, S., Ramesh B, Umamahesh K, Vijaya S. 2016. Probiotication of Tomato and Carrot Juices for Shelf-life. Enhancement using Micro-encapsulation. *Journal of Food Biosciences and Technology*. 6(2):13-22.
- Nagpal, R., Kumar, A. and Kumar M. 2012. Fortification and fermentation of fruit juices with probiotic lactobacilli. *Annals of Microbiology*. 62, 1573–1578.
- Nazzaro, F., Fratianni, F., Sada, A. and Orlando, P. 2008. Synbiotic potential of carrot juice supplemented with *Lactobacillus* spp. and inulin or fructooligosaccharides. *Journal of the Science of Food and Agriculture*. 88, 2271-2276
- Nuallkaekul S, Salmeron I, Charalamm popoulos D. 2011. Investigation of the factors influencing the survival of *Bifidobacterium longum* in model acidic solutions and fruit juices. *Food Chemistry*. 129, 1037-44.
- Nuallkaekul, S. and Charalampopoulos, D. 2011. Survival of *Lactobacillus plantarum* in model solutions and fruit juices. *Int. J. Food Microbiol*. 146, 111–117.
- Pakbin, B., Razavi, S.H., Mahmoudi, R. and Gajarbeygi, P. 2014. Producing probiotic peach juice. *Biotechnology Health Science*. 1, 1–5.
- Panda, S.H., and R. C. Ray. 2007. Lactic acid fermentation of  $\beta$ - carotene rich sweet potato (*Ipomoea batatas*L.) into lacto-juice,” *Plant Foods for Human Nutrition*. 62(2): 65–70.
- Patel, A.R. 2017. Probiotic fruit and vegetable juices-recent advances and future perspectives. *Int. Food Res. J*. 24, 1850–1857.
- Peerajan, S., Chaiyasut, C., Sirilun, S., Chaiyasut, K., Kesika, P. and Sivamaruthi, B.S. 2016. Enrichment of nutritional value of *Phyllanthusemblica* fruit juice using the probiotic bacterium, *Lactobacillus paracasei* HII01 mediated fermentation. *Food Science and Technology*. 36(1):116-123.
- Peres, C., Hernández-Mendoza, A. and Malcata, F.X. 2012. Review on fermented plant materials as carriers and sources of potentially probiotic lactic acid bacteria- With an emphasis on table olives. *Trends in Food Science and Technology*. 26, 31–42.
- Perricone, M.; Corbo, M.R.; Sinigaglia, M.; Speranza, B.; Bevilacqua, A. 2014. Viability of *Lactobacillus reuteri* in fruit juices. *J. Funct. Foods*. 10, 421–426.
- R. Di Cagno, R. Coda, M. De Angelis, and M. Gobbetti. 2013. Exploitation of vegetables and fruits through lactic acid

- fermentation. *Food Microbiology*. 33(1): 10.
- Rakin, M., Vukasinovic, M., Siler-Marinkovic, S. and Maksimovic, M. 2007. Contribution of lactic acid fermentation to improved nutritive quality vegetable juices enriched with brewer's yeast autolysate. *Food Chemistry*. 100, 599–602.
- Rolle R, Satin M. 2002. Basic requirements for the transfer of fermentation technologies to developing countries. *International Journal of Food Microbiology*. 75(3):181-187.
- Saarela, M., Alakomi, H.L.; Mättö, J.; Ahonen, A.M.; Tynkkynen, S. 2011. Acid tolerant mutants of *Bifidobacterium animalis* subsp. *lactis* with improved stability in fruit juice. *Food Sci. Technol*. 44, 1012–1018.
- Saarela, M., L. L'ahteenmäki, R. Crittenden, S. Salminen, and T. Mattila-Sandholm. 2002. Gut bacteria and health foods—the European perspective. *International Journal of Food Microbiology*. 78(1): 99–117.
- Saarela, M., Virkajarvi, I., Nohynek, L., Vaari, A. and Matto, J. 2006. Fibres as carriers for *Lactobacillus rhamnosus* during freeze-drying and storage in apple juice and chocolate-coated breakfast cereals. *International Journal of Food Microbiology*. 112, 171–178.
- Šárka Horáčková, Kristýna Rokytová, Kristina Bialasová, Iveta Klojdová and Marcela Sluková. 2018. Fruit juices with Probiotics – New type of Functional Foods. *Czech J. Food Sci*. 36(4): 00-5.
- Scheinbach, S., 1998. Probiotics: Functionality and commercial status, *Biotechnol. Adv*. 16, 581–608.
- Shah, N., and Prajapati J.B. 2013. Effect of carbon dioxide on sensory attributes, physico-chemical parameters and viability of Probiotic *L. helveticus* MTCC 5463 in fermented milk. *Journal of food Science and Technology*. 51 (12): 3886-3893.
- Shah, NP, Ding WK, Fallourd MJ and Leyer G. 2010. Improving the stability of probiotic bacteria in model fruit juices using vitamins and antioxidants. *Journal of Food Science*. 75(5): 278-82.
- Sheehan, V.M., Ross, P. and Fitzgerald, G.F. 2007. Assessing the acid tolerance and the technological robustness of probiotic cultures for fortification in fruit juices. *Innovative Food Science and Emerging Technologies*. 8: 279–284.
- Sima Khezri, Razzagh Mahmoudi and ParvinDehghan. 2018. Fig juice fortified with inulin and *L. delbrueckii*: A Promising Functional Food. *Applied Food Biotechnology*. 5(2): 97-106.
- Sivudu SN., Umamahesh K, Reddy OVS. 2014. A Comparative study on Probiotication of mixed watermelon and tomato juice by using probiotic strains of lactobacilli. *International Journal of Current Microbiology and Applied Sciences*. 3(11):977-984.
- Soccol, C.R., F.C. Prado, J.L. Parada. 2007. Technological process to produce a coconut fermented beverage with probiotic properties. *BR patent PI0703244-7* (in Portuguese).
- Sohail, A., Turner, M.S. Prabawati, E.K. Coombes, A.G.A. and Bhandari, B. 2012. Evaluation of *Lactobacillus rhamnosus* GG and *Lactobacillus acidophilus* NCFM encapsulated using a novel impinging aerosol method in fruit juice products. *International Journal of Food Microbiology*. 157, 162–166.
- Steinkraus KH., 1997. Classification of fermented foods: worldwide review of household fermentation techniques. *Food Control*. 8, 311-317.
- Tamang, J.P., 2009. Himalayan Fermented Foods: Microbiology, Nutrition and Ethnic Values. CRC Press, New Delhi, India.
- Tamime, A.Y., M. Saarela, A. Korslund Søndergaard, V.V. Mistry, N.P. Shah. 2005. Production and Maintenance of Viability of Probiotic Micro-Organisms in Dairy Products. In: *Probiotic Dairy Products*, A.Y. Tamime (Ed.), Blackwell Publishing, Oxford, UK. pp. 44–51.

- Tripathi, M.K., Giri, S.K. 2014. Probiotic functional foods: Survival of probiotics during processing and storage. *J. Funct. Foods*.9, 225–241.
- Tsen, J.H., Lin, Y.P. and King, V.A.E. 2004. Fermentation of banana media by using  $\kappa$ -carrageenan immobilized *Lactobacillus acidophilus*. *International Journal of Food Microbiology*. 91, 215–220.
- Tuorila, H., and Cardello, A. V. 2002. Consumer responses to an off-flavour in juice in the presence of specific health claims. *Food Quality and Preference*. 13: 561–569.
- Valero-Cases, E.; Frutos, M.J. 2017. Effect of Inulin on the Viability of *L. plantarum* during Storage and In Vitro Digestion and on Composition Parameters of Vegetable Fermented Juices. *Plant Foods Hum. Nutr.* 72, 161–167.
- Vasudha S., Mishra H. 2013. Non Dairy Probiotic Beverages. *International Food Research Journal*. 20(1):7-15.
- Vijaya Kumar, B., K. Umamahesh and O. V. S. Reddy. 2017. Isolation and identification of *Pediococcus pentosaceus* from cow milk's curd and its use in grape juices fermentation. *Journal of Advances in Biology and Biotechnology*. 13(4): 1-11.
- Wang, C.Y., Ng, C.C., Su, H., Tzeng, W.S., and Shyu, Y.T. 2009. Probiotic potential of noni juice fermented with lactic acid bacteria and bifidobacteria. *International Journal of Food Science and Nutrition*. 60(6): 98-106.
- Yoon, K.Y., Woodams, E. E. and Hang, Y. D. 2004. Probiotication of tomato juice by lactic acid bacteria. *Journal of Microbiology*. 42, 315–318.
- Yoon, K.Y., Woodams, E.E.; Hang, Y.D. 2005. Fermentation of beet juice by beneficial lactic acid bacteria. *LWT Food Sci. Technol.* 38, 73–75.
- Yoon, K.Y., Woodams, E.E.; Hang, Y.D. 2006. Production of probiotic cabbage juice by lactic acid bacteria. *Bioresour. Technol.* 97, 1427–1430.

**How to cite this article:**

Aditya Chaudhary. 2019. Probiotic Fruit and Vegetable Juices: Approach Towards a Healthy Gut. *Int.J.Curr.Microbiol.App.Sci*. 8(06): 1265-1279.  
doi: <https://doi.org/10.20546/ijcmas.2019.806.154>