

Original Research Article

<https://doi.org/10.20546/ijcmas.2019.807.160>

In-vitro* Multiple Shoots Production from Cormel Shoot Buds in *Gladiolus (Gladiolus hybrida)

Parul Devi, Pushpendra Kumar*, R.S. Sengar, M.K. Yadav,
Mukesh Kumar, S.K. Singh and Shilpy Singh

Department of Agricultural Biotechnology, Sardar Vallabhbhai Patel University of Agriculture
and Technology, Meerut, U.P. - 250110, India

*Corresponding author

ABSTRACT

Keywords

*Gladiolus
hybrida*,
in vitro
regeneration,
Growth
regulators,
2,4-D, BAP,

Article Info

Accepted:
12 June 2019
Available Online:
10 July 2019

Gladiolus (Gladiolus hybrida) is an important ornamental plant cultivated world over for its attractive spikes. The commercial cultivation of *Gladiolus* is based on natural multiplication of corms and cormels. *In vitro* propagation techniques, is superior to conventional propagation and produces disease free plants in huge quantities. *In vitro* regeneration of gladiolus cultivars, *Sylvia*, *White Prosperity* and *Amsterdam* was achieved using shoot bud of cormels as an explant. *Sylvia* variety recorded the best for callus induction and regeneration than other two varieties. The concentration of plant growth regulators affects the regeneration response differently. The highest callus induction rate of 76% explants was recorded on Murashige and Skoog (MS) medium supplemented with 2 mg^l⁻¹ 2,4-D (2,4-Dichlorophenoxyacetic acid), after 28 days. MS medium supplemented with BAP at 2.0 mg^l⁻¹ exhibited higher shoot proliferating efficiency, i.e shoots per explant in *Sylvia* variety.

Introduction

Gladiolus (Gladiolus hybrida) is a bulbous ornamental plant. It has a great commercial value in cut flower industry all over the world as well as in India due to its magnificent and colourful spikes. The genus *Gladiolus* belongs to the family Iridaceae. United States (Florida and California), Holland, Italy, France, Poland, Bulgaria, Brazil, India, Australia and Israel are the major gladiolus producing

countries. The major gladiolus producing states in the India are Uttar Pradesh, West Bengal, Odisha, Chhattisgarh, Haryana and Maharashtra. *Gladiolus* is also grown in states like Uttarakhand, Karnataka, Andhra Pradesh and Sikkim (Kadam *et al.*, 2014). Commercial production of corms and cormels of gladiolus is affected by *Fusarium* corm rot during storage. Besides, about 25 cormels produce from one mother corm in each season (Sinha and Roy, 2002). These cormels were not able

to be directly forced to flower, since gladiolus plant was naturally grown as biennial. Thus its commercial cultivation is limited by low rate of multiplication and does not fulfill the local demand of planting material which eventually affects the final cost of corms.

Therefore novel cultivars need to be rapidly mass multiplied by using modern *in vitro* technologies in order to fulfill the supply gap of huge demand of our local market which is of course not possible through conventional methods. Through modern technologies Mass propagation of corms and cormels such as tissue culture techniques have adopted at commercial level. Advanced countries are using highly sophisticated modern technologies for the commercial production of desired varieties in order to compete in the international markets. *In-vitro* plant tissue culture makes it possible to produce disease free and true to type planting material of Gladiolus.

In-vitro propagation techniques, prove significance especially for securing rapid multiplication of the novel cultivars. Although there are several research on *in vitro* propagation of gladiolus varieties using shoot bud, root, leaf and other different parts of plant as explant, and various plant growth regulators such as 2,4-D, IAA, NAA and BAP (Misra *et al.*, 1999; Pathania *et al.*, 2001; Kumari *et al.*, 2005 and Roy *et al.*, 2006). *In-vitro* micropropagation of gladiolus has been reported by using axillary buds (Begum *et al.*, 1995; Boonvanno *et al.*, 2000), shoot tip (Hussain *et al.*, 2001), cormels (Nagaraju *et al.*, 1995) and inflorescence axes (Ziv *et al.*, 2000). Successful protocols for *in vitro* corm formation (Dantu and Bhojwani, 1995; Sen *et al.*, 1995; Al-Juboory *et al.*, 1995), organogenesis and somatic embryogenesis (Remotti *et al.*, 1995; Kumar *et al.*, 2002) have been achieved also. However, there is a clear scope for further refinement through *in*

vitro culture methodology to acquire a higher number of shoots to complement traditional nursery methods in Gladiolus (Hussain *et al.*, 2001). Present study has been undertaken for standardization of plant growth regulators on MS medium supplemented with PGRs for *in vitro* differentiation and regeneration of three popular varieties of gladiolus.

Materials and Methods

Procurement and preparation of explants

The healthy corms of gladiolus cv. *Sylvia*, *White prosperity* and *Amsterdam* were obtained from Department of Horticulture of Sardar Vallabh Bhai Patel University of Agriculture and Technology (SVPUA&T), Modipuram, Meerut. Plants were raised in the field laboratory of Department of Agriculture Biotechnology, SVPUA&T, Meerut for collecting the cormels of gladiolus. These cormels were used for the research work. The outer scale of cormels was removed and shoot buds were cut with the help of surgical blade. Then buds were washed with 3-4 drops of Twin-20 (liquid detergent) along with sodium hypochlorite solution. Then dipped into 1% bavistin for 10 min followed 0.1 % HgCl₂ solution for 2 minutes. After each treatment, the bud were washed with sterile distilled water, 4 to 5 times. Buds were dried using the blotting paper before inoculated on the media.

Culture media

Media for callus induction

MS medium supplemented with various concentrations of plant growth regulators was used for the present study. Callus induction initiated when surface sterilized explants were inoculated on MS medium containing 2,4-D, NAA and IBA. Sucrose 3% (w/v) was added as carbon source. The pH was adjusted to 5.8 by 1N NaOH or 1N HCl. Agar 0.7% was

added as a gelling agent in the medium.

Media for shoot regeneration

After Callus initiation the explants were placed on MS medium supplemented with various concentrations of BAP, KIN for shoot regeneration. Sucrose 3% (w/v) was added as carbon source. The pH was adjusted to 5.8 by 1N NaOH or 1N HCl. Agar 0.7% was added as a gelling agent in the medium. All cultures were maintained at 25 ± 2°C with 16/8 h (light/dark) photoperiod.

Results and Discussion

In shoot bud culture, plantlets were regenerated via indirect organogenesis, In indirect approach, plantlets were regenerated from callus mass. Cormels shoot bud were cut and transferred into MS medium with varying hormonal concentrations of 2,4-D (0.5 to 4 mg^l⁻¹), NAA (0.5 to 4 mg^l⁻¹) and IBA (0.5 to 4 mg^l⁻¹). Callus induction was successfully observed in the shoot bud explants of *Sylvia* variety. In *Sylvia* variety excellent callus formation was observed while in *White Prosperity* and *Amsterdam* poor callus formation was observed (Table 1). After four

weeks of inoculation, *Sylvia* variety showed better callus formation as compared to other two varieties. Maximum mass of callus was recorded at the concentrations of auxin 2 mg^l⁻¹ 2,4-D followed by 2 mg^l⁻¹ IBA on full strength of MS medium. Kamo (1994) and Grewel *et al.*, (1995) also documented similar findings for diverse explant cultures in gladiolus.

Effect of PGRs on *in vitro* shoot differentiation from callus of gladiolus

Shoot regeneration from callus was done by transferring it to MS medium containing various concentrations of Plant Growth Regulators. Shoot induction was observed after 10-20 days of transfer on shoot development media (MS medium supplemented with BAP 1.0 mg^l⁻¹ and 2.0 mg^l⁻¹) (Fig 1.B). The result of mean value showed that among all the varieties studied, number of shoots was maximum with the concentration of BAP (1.0 mg^l⁻¹ and 2.0 mg^l⁻¹) in *Sylvia* variety. While *White Prosperity* and *Amsterdam* variety showed poor number of shoot regeneration (Fig 1. C, D; Table 2). Earlier studies also reported that cytokinin enhances shoot organogenesis (Remotti, 1995; Kumar *et al.*, 2002 ; Aslam *et al.*, 2012).

Table.1 MS basal medium supplemented with different concentration of plant growth regulators for callus induction in gladiolus cultivars, *Sylvia*, *White prosperity* and *Amsterdam*


S. No.	Treatments	Media	Varietal response to callus formation		
			<i>Sylvia</i>	<i>White prosperity</i>	<i>Amsterdam</i>
1	T ₁	MS+0.5 mg/l 2,4-D	+++	++	-
2	T ₂	MS+1.0 mg/l 2,4-D	+++	+	-
3	T ₃	MS+2.0 mg/l 2,4-D	++++	++	+
4	T ₄	MS+4.0 mg/l 2,4-D	++	+	++
5	T ₁	MS+0.5 mg/l NAA	+	-	-
6	T ₂	MS+1.0 mg/l NAA	+	-	-
7	T ₃	MS+2.0 mg/l NAA	++	-	-
8	T ₄	MS+4.0 mg/l NAA	+++	+++	+
9	T ₁	MS+0.5 mg/l IBA	+	-	-
10	T ₂	MS+1.0 mg/l IBA	+++	+	-
11	T ₃	MS+2.0 mg/l IBA	++++	++	+
12	T ₄	MS+4.0 mg/l IBA	++	+	++

Abbreviation use in this table: ++++ very good, +++ good, ++ poor, + very poor and -No callus

Table.2 MS basal medium supplemented with different concentration of cytokinins for regeneration from callus in gladiolus cultivars, *Sylvia*, *White prosperity* and *Amsterdam*

S. No.	Treatments	Media	Varietal response to micro shoots production (number)		
			Sylvia	White prosperity	Amsterdam
1	T ₁	MS+0.5 mg/l BAP	3.60 ± 0.24	2.11 ± 0.5	5.40 ± 0.12
2	T ₂	MS+1.0 mg/l BAP	9.00 ± 0.55	1.00 ± 0.25	4.70 ± 0.37
3	T ₃	MS+2.0 mg/l BAP	9.20 ± 0.86	2.30 ± 0.50	3.40 ± 0.81
4	T ₄	MS+4.0 mg/l BAP	4.60 ± 0.40	5.90 ± 0.21	3.11 ± 0.71
5	T ₁	MS+0.5 mg/l KIN	5.60 ± 0.75	10.90 ± 0.56	6.12 ± 0.87
6	T ₂	MS+1.0 mg/l KIN	4.20 ± 0.58	9.81 ± 0.67	4.23 ± 0.56
7	T ₃	MS+2.0 mg/l KIN	5.80 ± 0.73	7.23 ± 0.66	5.20 ± 0.57
8	T ₄	MS+4.0 mg/l KIN	6.40 ± 0.51	6.42 ± 0.23	6.60 ± 0.68
		Mean ± Std. error	6.05 ± 0.07	5.67 ± 1.27	4.85 ± 0.98

Fig.1 A: Callus induction from cormel shoot buds of gladiolus cv. *Sylvia* inoculated on full strength MS medium supplemented with 2,4-D (2.0 mg l⁻¹), twenty eight days after inoculation.
B: Shoot induction in eight week old callus of *gladiolus* cv. *Sylvia* inoculated on MS medium supplemented with BAP (1.0 mg l⁻¹ and 2.0 mg l⁻¹), ten-twenty days after inoculation
C: Shoot induction of gladiolus cv. *White Prosperity* inoculated on MS medium supplemented with BAP (1.0 mg l⁻¹ and 2.0 mg l⁻¹), ten-twenty days after inoculation.
D: Shoot induction of gladiolus cv. *Amsterdam* inoculated on MS medium supplemented with BAP (1.0 mg l⁻¹ and 2.0 mg l⁻¹), ten-twenty days after inoculation.



For gladiolus tissue culture experiments, mostly MS basal medium was used by different scientists (Hussey, 1977; Longan and Zettler, 1985; Kamo *et al.*, 1990; Dantu and Bhojwani, 1992). During the present study, basal MS medium was used throughout the experiment as this has been reported more responsive than other medium. Various types and concentrations of plant growth regulators in different combinations were supplemented into basal MS medium. During present investigation, response of three varieties of gladiolus were recorded on MS media supplemented with different auxins and cytokinins concentrations.

References

- Al-Juboory K.H., Shibli R.A., and Skiryin, R. (1995). Organogenesis and cormel production from callus culture of gladiolus cv. *Balady*. *Mu'tah. J. Res. Stud.*, 12: 143-160.
- Aslam, F., Habib, S. and Naz, S. (2012). Effect of different phytohormones on plant regeneration of *Amaryllis hippeastrum*. *Pakistan Journal of Science* 64: 54-58.
- Begum S. and Haddiuzaman S. (1995). *In vitro* rapid shoot proliferation and corm development in *Glaiolus grandiflorus* cv. *Red brand*. *Plant Tissue Cult.*, 5: 712.
- Boonvanno K. and Kanchanapoom K. (2000). *In-vitro* propagation of gladiolus. *Suranaree J. Sci. Technol.* 7: 25-29.
- Dantu P.K and Bhojwani S.S (1995). *In vitro* corm formation and field evolution of corm derived plants of gladiolus. *Sci. Hort.*, 61: 115–129.
- Grewel, M. S., Arora, J. S. and Gosal, S. S. (1995). Micropropagation of gladiolus through *in vitro* cormel production. *Plant Tissue Culture* 5: 27-33.
- Hussain I., Muhammad A., Rashid H. and Quraishi A. (2001). *In vitro* multiplication of gladiolus (*Gladiolus crassifolius*). *Plant Tissue Cult.* 11: 121-126.
- Hussain, S. C. T., Geetha, C. K., Rajeevan, P. K. and Valsalakumari, P. K. (1994). Plant regeneration from root derived callus in gladiolus. *Journal of Ornamental Horticulture* 2:46-50.
- Hussey G. (1977). *In vitro* propagation of gladiolus by precocious auxiliary shoot formation. *Scientia Horticulturae* 6:296.
- Kadam J.J, Agale R.C, Rite S.C, Pandav S.M. (2014). Exploration of fungicides and phytoextract against *Fusarium oxysporum* sp. Gladioli causing corm rot of gladiolus. *Discovery Agriculture*, 2(9): 61-64.
- Kamo K., Chen J. and Lawson R. (1990). The establishment of cell suspension cultures of gladiolus that regenerate plants *in vitro*. *Cell and Developmental Biology* 26:425-430.
- Kamo, K. (1994). Effect of phytohormones on plant regeneration from callus of gladiolus cultivar's Jenny Lee *in vitro* *Cell and Developmental Biology* 30: 26-31.
- Kumar A., Palni L.M.S. Sood A., Sharma M., Palni U.T. and Gupta A.K. (2002). Heat shock induced somatic embryogenesis in callus cultures of gladiolus in the presence of high sucrose. *Journal of Hort. Sci. and Biotech*, 77(1): 73-78
- Longan A. E. and Zettler F. (1985). Rapid *in vitro* propagation of virus indexed gladiolus. *Acta Horticulturae* 164:169-80.
- Misra S. and Singh R. (1999). *In vitro* propagation of gladiolus cv. 'American Beauty'. *J. Ornament. Hort.*, 2: 67-70.
- Nagaraju V. and Parthasarthy V.A. (1995). Effect of growth regulators on *in vitro* shoots of *Gladiolus hybridus*. *Folia Hort*, 7: 93-100.
- Pathania N.S., Misra R.L. and Raghava S.P.S. (2001). Precocious shoot proliferation

- and microcorm production in gladiolus through tissue culture. *Journal of Ornamental Horticulture*, 4: 69-73.
- Priyakumari I. and Sheela V. L. (2005). Micropropagation of gladiolus cv. 'Peach blossom' through enhanced release of auxiliary buds. *Journal of Tropical Agriculture* 43:47-50.
- Remotti P.C. and Loffler H.J.M. (1995). Callus induction and plant regeneration from gladiolus. *Plant Cell, Tissue and Organ Culture*, 42: 171-178.
- Roy S.K., Gangopadhyay G., Bandyopadhyay T., Modak B.K., Datta S. and Mukherjee K.K. (2006). Enhancement of *in vitro* micro corm production in gladiolus using alternative matrix. *African J. Biotechnol.*, 5: 1204-1209.
- Sen J. and Sen S. (1995). Two step bud culture technique for a high regeneration of gladiolus corm. *Sci. Hort.*, 64: 133-138.
- Sinha P. and Roy S.K. (2002). Plant regeneration through *in vitro* cormel formation from callus culture of *Gladiolus primulinus* Baker. *Plant Tiss. Cult*, 12: 139- 145.
- Ziv M. and Lilien-Kipnis H. (2000). Bud regeneration from inflorescence explants for rapid propagation of geophytes *in vitro*. *Plant Cell Rep.*, 19: 845-850.

How to cite this article:

Parul Devi, Pushpendra Kumar, R.S. Sengar, M.K. Yadav, Mukesh Kumar, S.K. Singh and Shilpy Singh. 2019. *In-vitro* Multiple Shoots Production from Cormel Shoot Buds in Gladiolus (*Gladiolus hybrida*). *Int.J.Curr.Microbiol.App.Sci.* 8(07): 1345-1350.
doi: <https://doi.org/10.20546/ijemas.2019.807.160>