

Original Research Article

<https://doi.org/10.20546/ijcmas.2020.907.216>

Development of Microbial Consortia of Nitrogen Fixing, Phosphate Solubilizing and Potash Mobilizing Bacteria for Optimizing Nutrient Supplementation to Soybean

J. R. Ghadge, D. R. Murumkar* and M. H. Shete

Department of Plant Pathology and Agril Microbiology, College of Agriculture,
Pune-411 005, Maharashtra, India

*Corresponding author

ABSTRACT

The present study was aimed with the formulation of suitable culture media for growth of nitrogen fixing, phosphate solubilizing and potash mobilizing bacteria in a consortium. Among different culture media, MS III media having Glucose (10 g l^{-1}), Mannitol (10 g l^{-1}), Ammonium sulphate (0.5 g l^{-1}) and Yeast extract (6 g l^{-1}) recorded maximum growth and microbial count of Rhizobium, PSB and KMB. These three beneficial microorganisms found compatible with each other when grown on MS III culture media. Furthermore, a field experiment was conducted to study the effect of seed inoculation of consortium of Rhizobium, PSB and KMB on growth parameters and yield of soybean. Among different inoculation treatments, seed inoculation with consortium of Rhizobium, PSB and KMB + 75% RDF was found to be the most effective as it recorded significantly highest germination (97.34%), shoot length (23.03 cm), root length (11.60 cm) and plant vigour index (3370.47) at 15 days after sowing, plant height (33.47 cm and 44.83 cm), root length (13.45 cm and 19.47 cm), dry weight of shoot ($7.65 \text{ g plant}^{-1}$ and $8.90 \text{ g plant}^{-1}$) and dry weight of root ($905.33 \text{ mg plant}^{-1}$ and $978 \text{ mg plant}^{-1}$) at flowering and harvest stage of the crop, number of branches (5.67 plant^{-1}), number of nodules (20.93 plant^{-1}), number of pods (55.23 plant^{-1}), 1000 seed weight (125.36 g), and seed yield (19.48 q ha^{-1}) of soybean and found statistically indistinguishable with the treatment of seed inoculation with consortium + 100% RDF for growth parameters and seed yield of soybean. The results indicated 25% saving of nitrogen, phosphorus and potassium dose of chemical fertilizers to soybean. Moreover, MS III culture medium proved effective with respect to population stability of individual strain and effectiveness of consortium of Rhizobium, PSB and KMB on growth and yield of soybean.

Keywords

Consortium,
Rhizobium, PSB,
KMB, Soybean

Article Info

Accepted:

17 June 2020

Available Online:

10 July 2020

Introduction

Microorganisms usually need different types of culture media for their growth under *in-vitro* condition. Yeast extract mannitol agar

media, Pikovskaya's media and Alexandrov media are suitable for individual growth of *Rhizobium*, phosphate solubilizing bacteria and potash mobilizing bacteria, respectively. But it needs to be essential to formulate such

a culture medium which found suitable for growth of these three beneficial microorganisms in a consortium. It is well known that phosphate solubilizing bacteria and *Rhizobium* have synergistic effect on legume crops. Development of consortia containing one strain of *Rhizobium*, PSB and PGPR has been attempted (Bansal, 2015).

The present study was undertaken for formulation of suitable culture media for growth of nitrogen fixing, phosphate solubilizing and potash mobilizing bacteria in a consortium and its application as seed inoculation under glasshouse experiment with soybean crop.

Materials and Methods

Isolation of *Rhizobium* from root nodules of soybean

The healthy, unbroken, firm and pink nodules from soybean roots were selected for isolation of *Rhizobium* by using yeast extract mannitol agar (YEMA) media as described by Rajendran *et al.*, (2008).

Nitrogen fixing ability of the rhizobial isolate

The 48 hour old culture of freshly isolated *Rhizobium* strain was inoculated to 5 ml of yeast extract mannitol medium. It was incubated for 48 hours. One ml of this broth was inoculated to 50 ml yeast extract mannitol medium. Then it was incubated for 15 days. Ten ml of this culture was used for N estimation by following the standard procedure of Microkjeldhal technique (Reis *et al.*, 1994). The formula for N₂ estimation is:

$$N_2 \text{ (mg/g)} = \frac{\text{ml of H}_2\text{SO}_4 \text{ in the sample} \times \text{Normality of H}_2\text{SO}_4 \times 14.01}{\text{Weight of the sample (carbon used in g)}}$$

Biochemical and physiological characterization of rhizobial isolate

Pure culture of the isolate was made and then subjected to Gram reaction. The Gram negative isolates were further subjected to biochemical tests including catalase, oxidase, gelatin hydrolysis, indole tests and growth on different carbon sources for confirmation. The biochemical characterization of the isolates was carried out as per the procedures outlined by Cappuccino and Sherman (1987) in their 10th edition of Microbiology: A Laboratory Manual.

Isolation of phosphate solublizing bacteria (PSB) from rhizosphere soil of soybean

The isolation of phosphate solublizing bacteria on Pikovskaya's medium was carried out by serial dilution of soil and agar plating method (Aneja, 2003). The formation of clear zone of P-solublization around the colonies grown on Pikovskaya's medium were selected, purified, subcultured and maintained on the slants of Pikovskaya's agar for further use.

Phosphate solublizing ability of the bacterial isolates

The ability of the bacterial isolates to solubilize insoluble inorganic phosphate was tested by spotting 10 µl overnight cultures on Pikovskaya's agar plates and incubating at 28-30°C for 2-3 days. The isolates which showed clear zone of solublization of tricalcium phosphate (TCP) around the colony were noted as phosphate solubilizers. The diameter of the zone of TCP solublization was measured and expressed in millimeters. The bacterial isolates positive for P solublization on Pikovskaya's agar medium were subjected to quantification of Pi released from TCP in broth medium.

Biochemical characterization of the PSB isolate

The biochemical characterization of the isolate was carried out as per the procedures outlined by Cappuccino and Sherman in their 10th edition of *Microbiology: A Laboratory Manual*. Catalase test, Oxidase test, Indole production test, Methyl red test, Voges-Proskauer (VP) test, Urea hydrolysis, Nitrate reduction test, Gelatin hydrolysis test, Starch hydrolysis, Casein hydrolysis and H₂S production test were performed.

Isolation of potash mobilizing bacteria from rhizospheric soil of soybean

One gram of rhizosphere soil was mixed thoroughly in 100 ml sterile water and was processed following serial dilution agar plate technique (Aneja, 2003). A suitable dilutions (10⁻⁵ and 10⁻⁶) of both rhizosphere and rhizoplane solutions were plated on Alexandrov medium (Hu *et al.*, 2006). The plates were incubated at room temperature (30±1°C) for 3 days and the colonies exhibiting clear zones of solubilization of muscovite mica were selected purified, subcultured and maintained on the slants of Alexandrov medium for further use.

Quantitative estimation of 'K' solubilization

The isolates showing zone of solubilization on Alexandrov agar medium were further examined for their ability to release K from broth media. The amount of K released from muscovite mica in the broth by the isolates was studied at 7, 15 and 20 days after incubation (DAI) under laboratory condition (Parmar *et al.*, 2016).

Biochemical characterization of KMB isolate

The biochemical characterizations of the KMB isolate was carried out as per the

procedures outlined by Bergey's Manual of Systematic Bacteriology 9th Edition (1993).

Sugar utilization, Methyl red test, Voges-Proskauer (VP) test, Urea hydrolysis, Nitrate reduction test, Gelatin hydrolysis test, catalase test, starch hydrolysis, Casein hydrolysis and H₂S production test were performed.

Selection of culture medium

The culture media (MS I, MS II, MS III, MS IV and MS V) of various compositions were formulated as described by Shete *et al.*, (2019) and screened for growth of nitrogen fixing, phosphate solubilizing and potash mobilizing bacteria in broth by using various carbon sources like glucose, sucrose and nitrogen sources like ammonium sulphate and yeast extract in different concentrations along with different micronutrients (Table 1). The pH of all culture media was maintained in the range of 6.9 to 7.1

In vitro studies

Broth of each culture media *viz.*, MS I, MS II, MS III, MS IV and MS V were inoculated with efficient strains of *Rhizobium*, PSB and Potash mobilizing bacteria separately as well as in consortia and kept for incubation at 28±2°C for 5 days.

The cfu count of *Rhizobium*, PSB and potash mobilizing bacteria was recorded after incubation period of 5 days by using direct plate count technique. Before development of consortium, all strains were examined *in vitro* for their compatibility on selective medium by cross streak method (Ganesan and Gnanamanickam, 1987).

Observations on growth and cfu count of *Rhizobium*, PSB and Potash mobilizing bacteria in each culture media were recorded.

Preparation of consortium of *Rhizobium*, PSB and KMB on a selective medium

Inoculum of *Bradyrhizobium japonicum*, *Bacillus subtilis* and *Frateuria aurantia* was prepared in selective medium MS III (Shete *et al.*, 2019). The media was inoculated in 500 ml conical flask containing 150 ml medium and incubated at $28 \pm 2^\circ\text{C}$ under shaking at 100-150 rpm for three days to give an optical density of 0.5 recorded at 535 nm. Lignite powder used as carrier was sterilized at 121°C and 1.04 kg/cm^2 pressure for one hour and inoculated with broth cultures of *Bradyrhizobium japonicum*, *Bacillus subtilis* and *Frateuria aurantia* (100 ml per 500 g of lignite powder). Lignite powder based inoculum was incubated at $28 \pm 2^\circ\text{C}$ for three days by adding 10% sugar solution to increase the population of respective microbe. Inoculum of *Bradyrhizobium japonicum*, *Bacillus subtilis* and *Frateuria aurantia* having cfu of 2×10^7 per gram of lignite powder were applied to soybean as seed coating.

Field experiment

A field experiment was conducted during *kharif*, 2019 in the field at College of Agriculture, Pune to study the effect of seed inoculation with consortium of *Rhizobium*, PSB and KMB on growth parameters, nutrient uptake and yield of soybean. The soybean variety *Phule Sangam* was used as a test crop. The experiment was laid out in randomized block design with three replications and nine treatments.

Treatment details

The soybean seeds were treated before sowing as follows:

T₁: Consortium of *Rhizobium*, PSB and KMB
T₂: Consortium of *Rhizobium*, PSB and KMB

+ 100% RDF

T₃: Consortium of *Rhizobium*, PSB and KMB + 75% RDF

T₄: Consortium of *Rhizobium*, PSB and KMB + 50% RDF

T₅: *Rhizobium* + 75% recommended N + 100% recommended P₂O₅ and K₂O

T₆: PSB + 75% recommended P₂O₅ + 100% recommended N and K₂O

T₇: KMB + 75% recommended K₂O + 100% recommended N and P₂O₅

T₈: 100% RDF

T₉: Absolute control

Observations

The observations on germination (%), shoot length (cm), root length (cm) and plant vigour index at 15 days after sowing, plant height (cm), root length (cm), dry weight of shoot (g plant^{-1}) and dry weight of root (mg plant^{-1}) at flowering and harvest stage of the crop, number of branches, number of nodules and number of pods per plant, 1000 seed weight, NPK uptake (kg ha^{-1}) and seed yield (q ha^{-1}) of soybean were recorded. Plant vigour index was computed at 15 days after sowing using the formula: Plant vigour index = Germination % x [shoot length (cm) + root length (cm)]. Nitrogen content of plant was estimated by following Modified Kjeldahl's process and accordingly N uptake (kg ha^{-1}) was estimated as $\text{N\%} \times \text{total dry matter yield (kg ha}^{-1})/100$.

Microbial count of *Rhizobium*, PSB and KMB at flowering stage of soybean

Fresh root nodules of soybean at flowering stage were analyzed for rhizobial population on yeast extract mannitol agar media as described by Rajendran *et al.*, (2008). Moreover, rhizospheric soil samples at flowering stage of soybean were analyzed for microbial population of phosphate solubilizing bacteria (PSB) and potash mobilizing bacteria (KMB) using serial

dilution of soil and agar plating method (Aneja, 2003). The PSB and KMB population was enumerated on Pikovskaya's media and Alexandrov's agar media, respectively, at 10^6 dilutions. The plates were incubated at 28 ± 2 °C temperature for 72 hours and colonies were counted. The population was expressed as cfu g^{-1} soil.

Statistical Analysis

The data recorded on various parameters was subjected to statistical analysis by following standard method of analysis of variance. The level of significance used in 'F' and 't' tests was $P = 0.05$. Critical difference (CD) values were calculated where the 'F' test was found significant (Panse and Sukhatme, 1985).

Results and Discussion

Isolation of nitrogen fixing, phosphate solubilizing and potash mobilizing bacteria

The isolation of *Rhizobium* from root nodule of soybean (var. *Phule Sangam*) was done using yeast extract mannitol agar medium. The isolation procedure was carried out for all the three samples and three isolates were obtained as RH-I, RH-II and RH-III. Moreover, isolation of phosphate solubilizing bacteria on Pikovskaya's medium was carried out by serial dilution of soil and agar plating method (Aneja, 2003). The isolation procedure was carried out for all the three rhizosphere soil samples and the plates were observed for the appearance of bacterial colony showing clear zone of solubilization of tricalcium phosphate purified (TCP) on Pikovskaya's medium. Three isolates were obtained as P-I, P-II and P-III. Furthermore, isolation of potash mobilizing bacteria was carried out on Alexandrov medium. The isolation procedure was carried out for all the three rhizosphere soil samples on Alexandrov medium (Hu *et al.*, 2006). The plates were

observed for the appearance of bacterial colony showing clear zone of solubilization of insoluble potassium bearing mineral (mica). Three isolates were obtained as K-I, K-II and K-III.

Nitrogen fixing ability of *Rhizobium* isolate

All the three *Rhizobium* isolates of soybean alongwith MPKV strain (*Bradyrhizobium japonicum*) were subjected to know the nitrogen fixation by Microkjeldhal method (Table 2). The isolate RH-1 fixed highest amount of nitrogen (148.65 μg of nitrogen/mg of carbon used). This was followed by MPKV strain, RH-II and RH-III isolate (145.78, 129.76 and 121.84 μg of nitrogen/mg of carbon used, respectively). The results of the present investigation are in agreement with results of Hema and Savalgi (2017) who reported that isolate from maize GdM5 fixed about 142 μg of nitrogen/mg of carbon used.

Phosphate solubilizing ability of the PSB isolates

All the three PSB isolates alongwith MPKV strain (*Bacillus subtilis*) were tested for their ability to solubilize inorganic phosphate both qualitatively and quantitatively and their results are presented in Table 3. Quick analysis of P-solubilization was carried out on Pikovskaya's agar medium. All the three isolates were able to form zone of P-solubilization on the medium. The diameter of the zone of P-solubilization ranged from 3-6 mm in different isolates.

Quantitative estimation of Pi released from TCP for bacterial isolates

The amount of Pi released from tri-calcium phosphate by the PSB isolates alongwith MPKV strain (*Bacillus subtilis*) in Pikovskaya's broth was estimated at 10 days after inoculation. The amount of Pi released

from TCP by the isolates at 10 DAI ranged from 11.43 to 29.38 per cent (Table 3). The isolate P-I recorded highest P-solubilization (29.38%) than the other isolates tested.

Decrease in pH of medium during phosphate solubilization

The decrease in pH of TCP broth from initially adjusted pH of 7.0 was also noted at 10 days after inoculation. The maximum reduction in pH of the medium i.e. pH 3.48 was recorded by P-I isolate followed by MPKV strain (*Bacillus subtilis*), P-II and P-III isolates (3.50, 4.09 and 4.11, respectively) (Table 3). The decrease in pH of the medium with the amount of Pi released had positive correlation.

Quantitative estimation of 'K' solubilisation of the KMB isolates

The isolates showing zone of solubilization on Alexandrov agar medium were further examined for their ability to release 'K' from broth media. The amount of 'K' released from muscovite mica in the broth by the isolates alongwith MPKV strain (*Frateruria aurantia*) were studied at 7, 15 and 20 days after incubation (DAI) in lab condition and found in the range of 6.49 to 40.81 $\mu\text{g ml}^{-1}$ (Table 4). The results indicated that the amount of released 'K' increased as the days of incubation increases and the highest amount of 'K' present at 20 DAI. The maximum solubilization of muscovite mica was observed in K-I isolate (40.81 $\mu\text{g ml}^{-1}$) followed by MPKV strain (*Frateruria aurantia*) (39.25 $\mu\text{g ml}^{-1}$) at 20 DAI. The results of the present investigation are in agreement with the results of Parmar *et al.*, (2016) who isolated 25 potassium solubilizing bacterial isolates from the rhizosphere of maize from various areas of Navsari district and tested quantitative estimation of 'K' solubilisation of the highly efficient KMB

isolates. He further reported the amount of 'K' released from muscovite mica in the broth by the isolates in the range of 1.89 to 46.52 $\mu\text{g ml}^{-1}$.

On the basis of nitrogen fixing, phosphate solubilising and potash mobilizing ability, highly efficient nitrogen fixing *Rhizobium* isolate (RH-I), phosphate solubilising isolate (P-I) and potash mobilizing isolate (K-I) were further tested for different biochemical characterization.

Biochemical characterization of Rhizobium, PSB and KMB isolate

The highly efficient nitrogen fixing rhizobial isolate (RH-I) was tested for different biochemical characters *viz.*, gram staining, motility test, gelatin hydrolysis, catalase test, oxidase test, indole production test, starch hydrolysis, H₂S production, Voges- Proskauer test and growth on different carbon sources (Table 5). The cells of nitrogen fixing rhizobial isolate were rod shape, motile and gram negative in reaction. The nitrogen fixing rhizobial isolate was positive for catalase test, oxidase test, indole production test, starch hydrolysis, H₂S production and Voges-Proskauer test but was negative for gelatin hydrolysis. Glucose, sucrose and mannitol were used as a sole carbon source for growth by the nitrogen fixing rhizobial isolate. Based on biochemical and physiological characterization, the nitrogen fixing rhizobial isolate was identified as *Bradyrhizobium japonicum*. The results of the present investigation are in conformity with results of Jadhav (2013) who isolated rhizobia from root nodule of soybean cultivated in Latur area and further characterized these isolates biochemically for specific characters of *Bradyrhizobium japonicum* according to Burgey's Manual of Systematic Bacteriology. All the isolates were positive for most of characters specific for *Bradyrhizobium*

japonicum. Further all isolates tested negative for gelatin hydrolysis.

The highly efficient phosphate solubilizing bacterial isolate (P-I) was tested for different biochemical characters *viz.*, gram staining, motility test, gelatin hydrolysis, catalase test, oxidase test, starch hydrolysis, H₂S production and Voges- Proskauer test (Table 5). The cells of phosphate solubilizing bacterial isolate were rod shape, motile and gram positive in reaction.

The phosphate solubilizing bacterial isolate was positive for gelatin hydrolysis, catalase test, starch hydrolysis and Voges- Proskauer test but was negative for oxidase test and H₂S production. Based on biochemical and physiological characterization (Claus and Berkeley, 1986), the phosphate solubilizing bacterial isolate was identified as *Bacillus subtilis*.

The highly efficient potash mobilizing

bacterial isolate (K-I) was tested for different biochemical characters *viz.*, gram staining, motility test, methyl red test, Voges-Proskauer (VP) test, urea hydrolysis, nitrate reduction test, gelatine hydrolysis test, catalase test, starch hydrolysis, casein hydrolysis, H₂S production test and growth on different carbon sources (Table 5). The potash mobilizing bacterial isolate was rod shape, motile and gram negative in reaction. The potash mobilizing bacterial isolate was positive for gelatin hydrolysis, catalase test, starch hydrolysis, urea hydrolysis, casein hydrolysis test, nitrate reduction test and methyl red test but was negative for H₂S production and Voges- Proskauer test. Sucrose, mannitol and maltose were used as a sole carbon source for growth by the potash mobilizing bacterial isolates. Based on biochemical and physiological characterization (Parmar *et al.*, 2016), the potash mobilizing bacterial isolate was identified as *Frateuria aurantia*.

Table.1 Composition of culture media for consortia of *Rhizobium*, PSB and KMB

S.N.	Chemicals	Composition of culture media (g)				
		MS I	MS II	MS III	MS IV	MS V
1	Glucose	5	15.	10	0	20
2	Mannitol	15	5	10	20	0
3	Tri calcium phosphate	5	5	5	5	5
4	Ammonium sulphate	0.1	0.3	0.5	0.7	0.9
5	Yeast extract	2	4	6	8	10
6	Magnesium sulphate	0.1	0.1	0.1	0.1	0.1
7	Potassium chloride	0.2	0.2	0.2	0.2	0.2
8	Manganese sulphate	0.001	0.001	0.001	0.001	0.001
9	Ferrous sulphate	0.1	0.1	0.1	0.1	0.1
10	Calcium carbonate	2	2	2	2	2
11	Potassium alumino silicates	2	2	2	2	2
12	Dipotassium hydrogen orthophosphate	0.1	0.1	0.1	0.1	0.1
13	Sodium chloride	0.1	0.1	0.1	0.1	0.1
	pH	6.9	7.1	7.2	6.9	7.1
	Distilled water	1lit	1lit	1lit	1lit	1lit

Table.2 Nitrogen fixing ability of *Rhizobium* isolate of soybean by Microkjeldhal method

Sr. No.	Rhizobium isolate	Nitrogen fixing ability (µg of Nitrogen/mg of Carbon)
1.	RH-I	148.65
2.	RH-II	129.76
3.	RH-III	121.84
4.	MPKV strain (<i>Bradyrhizobium japonicum</i>)	145.78

Table.3 Zone of P solubilization on Pikovskaya's agar and per cent Pi released from TCP broth by the PSB isolates

Sr. No.	PSB Isolate	Zone of P solubilization on TCP (mm)	% Pi released from TCP after 10 days	Decrease in pH of medium (from initial pH 7.0) after 10 days
1	P-I	6.0	29.38	3.48
2	P-II	5.0	13.23	4.09
3	P-III	3.0	11.43	4.11
4.	MPKV strain (<i>Bacillus subtilis</i>)	6.0	28.35	3.50

Table.4 Solubilization of muscovite mica by the KMB isolates

Sr. No.	KMB isolate	7 DAI (µg ml ⁻¹)	15 DAI (µg ml ⁻¹)	20 DAI ((µg ml ⁻¹)
1.	K-I	24.56	37.26	40.81
2.	K-II	12.50	21.58	33.75
3.	K-III	6.49	19.60	30.42
4.	MPKV strain (<i>Frateuria aurantia</i>)	23.25	35.10	39.25

Table.5 Selective biochemical tests of nitrogen fixing, phosphate solubilizing and potash mobilizing bacterial isolate

Sr. No.	Biochemical tests	Rhizobium isolate (RH-I)	PSB isolate (P-I)	KMB isolate (K-I)
1.	Cell shape	Rod shape	Rod shape	Rod shape
2.	Gram reaction	Gram negative	Gram positive	Gram negative
3.	Motility	+	+	+
4.	Gelatin hydrolysis	-	+	+
5.	Catalase test	+	+	+
6.	Oxidase test	+	-	
7.	Indole production test	+		
8.	Starch hydrolysis	+	+	+
9.	H ₂ S production	+	-	-
10.	Voges- Proskauer test	+	+	-
11.	Urea hydrolysis			+
12.	Caesin hydrolysis test			+
13.	Nitrate reduction test			+
14.	Methyl red test			+
15.	Growth on carbon sources			
	a) Glucose	+		
	b) Sucrose	+		+
	c) Mannitol	+		+
	d) Maltose			+

Table.6 Growth of *Rhizobium*, PSB and KMB on different culture media

Sr. No.	Culture media	<i>Rhizobium</i>	PSB	KMB
1.	MS I	+	+	+
2.	MS II	++	+	-
3.	MS III	+++	+++	+++
4.	MS IV	+	-	-
5.	MS V	-	+	+

Table.7 Microbial count of *Rhizobium*, PSB and KMB in a consortium on different culture media

Sr. No.	Culture media	<i>Rhizobium</i> (cfu g ⁻¹)	PSB (cfu g ⁻¹)	KMB (cfu g ⁻¹)
1.	MS I	1 x 10 ³	1 x 10 ³	1 x 10 ³
2.	MS II	1 x 10 ⁵	1 x 10 ³	-
3.	MS III	12 x 10 ⁷	7 x 10 ⁷	9 x 10 ⁷
4.	MS IV	1 x 10 ³	-	1 x 10 ⁷
5.	MS V	-	1 x 10 ³	1 x 10 ⁷

Table.8 Inoculation effect of consortium of *Rhizobium*, PSB and KMB on growth parameters of soybean

Tr No	Treatment details	Germination (%)	Plant vigour index	Plant height (cm)		Root length (cm)		Dry weight of shoot (g plant ⁻¹)	
				Flowering	Harvest	Flowering	Harvest	Flowering	Harvest
T ₁	Consortium of <i>Rhizobium</i> , PSB and KMB	93.21	2840.5 2	31.47	42.60	12.02	16.62	6.49	7.82
T ₂	Consortium of <i>Rhizobium</i> , PSB and KMB + 100% RDF	96.60	3132.8 2	32.50	43.72	12.62	18.27	7.29	8.41
T ₃	Consortium of <i>Rhizobium</i> , PSB and KMB + 75% RDF	97.34	3370.4 7	33.47	44.83	13.45	19.47	7.65	8.90
T ₄	Consortium of <i>Rhizobium</i> , PSB and KMB + 50% RDF	95.06	2957.8 5	31.53	42.67	12.10	16.80	6.83	8.02
T ₅	<i>Rhizobium</i> + 75% recommended N + 100% recommended P ₂ O ₅ & K ₂ O	92.89	2867.9 0	31.37	41.87	11.90	16.42	5.95	7.22
T ₆	PSB + 75% recommended P ₂ O ₅ + 100% recommended N and K ₂ O	91.35	2776.7 7	30.87	41.58	11.75	16.28	5.62	6.97
T ₇	KMB + 75% recommended K ₂ O + 100% recommended N and P ₂ O ₅	91.00	2665.2 5	30.78	41.22	11.60	15.87	5.50	6.48
T ₈	100% RDF	91.76	2769.1 4	32.20	42.87	12.25	17.93	6.85	8.14
T ₉	Uninoculated control	85.18	2257.6 4	27.87	39.33	9.20	13.38	3.95	5.49
	S.E.	2.12	82.47	0.34	0.42	0.29	0.47	0.19	0.21
	C.D.at 5%	6.35	247.23	1.02	1.27	0.85	1.40	0.58	0.64
	C.V.	3.96	5.01	1.88	1.73	4.16	4.81	5.34	4.96

PSB = Phosphate solubilizing bacteria,

KMB = Potash mobilizing bacteria

Table.9 Inoculation effect of consortium of *Rhizobium*, PSB and KMB on growth and yield attributing characters of soybean

Tr. No	Treatment details	Dry weight of root (g plant ⁻¹)		Number of branches plant ⁻¹	Number of nodules plant ⁻¹	Number of pods plant ⁻¹	1000 seed weight (g)	Seed yield (q ha ⁻¹)
		Flowering	Harvest					
T ₁	Consortium of Rhizobium, PSB and KMB	803.33	879.67	7.00	18.20	49.97	120.00	17.04
T ₂	Consortium of Rhizobium, PSB and KMB + 100% RDF	850.00	918.33	7.13	19.85	53.20	123.04	18.44
T ₃	Consortium of Rhizobium, PSB and KMB + 75% RDF	905.33	978.00	7.67	20.93	55.23	125.36	19.48
T ₄	Consortium of Rhizobium, PSB and KMB + 50% RDF	808.33	885.67	7.03	18.33	50.27	120.18	17.18
T ₅	Rhizobium + 75% recommended N + 100% recommended P ₂ O ₅ & K ₂ O	788.67	853.00	6.97	18.13	48.87	119.76	16.62
T ₆	PSB + 75% recommended P ₂ O ₅ + 100% recommended N and K ₂ O	767.00	835.00	6.70	17.13	48.20	118.89	16.33
T ₇	KMB + 75% recommended K ₂ O +100% recommended N and P ₂ O ₅	742.33	829.67	6.63	16.83	48.00	117.27	16.21
T ₈	100% RDF	812.33	895.33	7.07	14.37	51.30	121.86	17.35
T ₉	Uninoculated control	377.00	412.00	5.60	9.77	30.33	109.74	14.62
	S.E.	25.97	25.05	0.19	0.37	0.79	0.83	0.70
	C.D.at 5%	77.85	75.10	0.58	1.10	2.37	2.50	2.09
	C.V.	5.91	5.22	6.88	3.74	2.83	1.21	7.08

Growth of Rhizobium, PSB and KMB on different culture media

Broth of each culture media viz., M I, M II, M III, M IV and M V were inoculated with efficient strains of *Rhizobium*, PSB and Potash mobilizing bacteria separately as well as in consortia and kept for incubation at $28 \pm 2^\circ\text{C}$ for 5 days. The data presented in Table 6 revealed that the maximum growth of *Rhizobium*, PSB and potash mobilizing bacteria was found on MS III culture media. Moreover, *Rhizobium*, PSB and KMB were found to be compatible with each other on MS III culture media (Fig. 2). Singh *et al.*, (2014) reported maximum growth of rhizobia in media containing 12.5 g l^{-1} sucrose at 29.4°C for 7 days. Further, Kucuk *et al.*, (2006) reported that *Rhizobium* strains were able to utilize glucose and sucrose more efficiently than normal YEM medium. Moreover, Sagervanishi *et al.*, (2014) studied the effect of different nitrogen sources viz., ammonium sulphate, casein, sodium nitrate and urea and found best optimized source was ammonium sulphate for the maximum 'P' solubilisation. Furthermore, Sugumaran and Janarthanam (2007) reported that *B. mucilaginosus* isolated from soil, rock and mineral samples recorded 4.29 mg l^{-1} release of potassium in media supplemented with muscovite mica. Results of the present investigation are in agreement with results of these researchers.

Microbial count of Rhizobium, PSB and KMB in a consortium on different culture media

The data on microbial count of *Rhizobium*, PSB and potash mobilizing bacteria is presented in Table 7. Among all the culture media, MS III culture medium recorded maximum count of *Rhizobium*, PSB and KMB (12×10^7 , 7×10^7 and 9×10^7 cfu g^{-1} , respectively).

Preparation of consortium of Rhizobium, PSB and KMB

Inoculum of *Rhizobium* (*Bradyrhizobium japonicum*), PSB (*Bacillus subtilis*) and KMB (*Frateuria aurantia*) was prepared in a selective medium MS III. The media was inoculated in 500 ml conical flask containing 150 ml medium and incubated at $28 \pm 2^\circ\text{C}$ under shaking at 100-150 rpm for three days to give an optical density of 0.5 recorded at 535 nm. Lignite powder used as carrier was sterilized at 121°C and 1.04 kg/cm^2 pressure for one hour and inoculated with broth cultures of *Bradyrhizobium japonicum*, *Bacillus subtilis* and *Frateuria aurantia* (100 ml in 500 g lignite powder). Lignite powder based inoculum was incubated at $28 \pm 2^\circ\text{C}$ for three days by adding 10% sugar solution to increase the population of the respective microbes. The inoculum of *Bradyrhizobium japonicum*, *Bacillus subtilis* and *Frateuria aurantia* having 2×10^7 cfu g^{-1} of lignite powder was applied to soybean as seed coating.

Inoculation effect of consortium of Rhizobium, PSB and KMB on growth and yield of soybean

The results in respect of growth and yield attributing characters of soybean are presented in Table 8 and 9. The results of the present investigation revealed that among the different inoculation treatments, T₃ i.e. seed inoculation with consortium of *Rhizobium*, PSB and KMB + 75% RDF was found to be the most effective as it recorded significantly highest germination (97.34%), plant vigour index (3370.47) at 15 days after sowing, plant height (33.47 cm and 44.83 cm), root length (13.45 cm and 19.47 cm), dry weight of shoot ($7.65 \text{ g plant}^{-1}$ and $8.90 \text{ g plant}^{-1}$) and dry weight of root ($905.33 \text{ mg plant}^{-1}$ and $978 \text{ mg plant}^{-1}$) at flowering and harvest stage of the crop, number of branches (5.67 plant^{-1}),

number of nodules (20.93 plant⁻¹), number of pods (55.23 plant⁻¹), 1000 seed weight (125.36 g) and seed yield (19.48 q ha⁻¹) of soybean, however it was statistically at par with the treatment T₂ i.e. seed inoculation with consortium + 100% RDF for growth parameters and seed yield of soybean. Bansal R. K. (2009) reported that presowing inoculation of mungbean seeds with different inoculants (*Rhizobium*, PGPR and PSB) alone or in combination, significantly increased the plant height, root length, dry matter production, number of nodules/plant, 1000 seed weight, nutrient uptake and seed yield over uninoculated control. Moreover, Qureshi *et al.*, (2011), Argaw (2012) and Tarafder *et al.*, (2016) reported increased growth parameters, nutrient uptake and seed yield in different legume crops due to seed inoculation of *Rhizobium*, PGPR and PSB alone or in combination. Results of the present investigation are in agreement with results of these researchers.

In conclusion MS III culture medium proved effective with respect to population stability of individual strain and effectiveness of consortium of *Rhizobium*, PSB and KMB on growth and yield of soybean. Moreover, from the present investigation it can be concluded that seed inoculation with consortium of *Rhizobium*, PSB and KMB + 75% RDF was found to be the most beneficial for getting higher seed yield of soybean with 25% saving of nitrogen, phosphorus and potassium dose of chemical fertilizers to soybean.

Acknowledgement

The authors express their sincere thanks and gratitude to the Head, Department of Plant Pathology and Agricultural Microbiology, College of Agriculture, Pune for extending research facilities and financial support for conducting the present study.

References

- Aneja, K.R. 2003. Experiments in Microbiology, Plant Pathology and Biotechnology. New Age International Publishers, New Delhi, India. pp. 1-607.
- Argaw A. 2012 Evaluation of Co-inoculation of *Bradyrhizobium japonicum* and Phosphate Solubilizing *Pseudomonas spp.* Effect on Soybean (*Glycine max* L. (Merr.)) in Assossa Area. *J. Agr. Sci. Tech.* 14: 213-224.
- Bansal, R. K. 2009. Synergistic effect of *Rhizobium*, PSB and PGPR on nodulation and grain yield of mung bean. *J. food legumes.* 22(1): 37-39.
- Bansal R K. 2015. Synergistic effect of *Rhizobium*, PSB and PGPR on nodulation and grain yield of mungbean. *International J. Agril. Biol.* 15: 55-68.
- Cappuccino, J.G. and Sherman, N. 1987. Microbiology A Laboratory Manual. 2nd Ed. The Benjamin/Cummins Publishing Co., USA., 458p.
- Claus, D. and Berkeley, R.C.W. 1986. Genus *Bacillus* cohn 1872. In: Bergy's Manual of Systematic Bacteriology, (Sneath, P.H.A., ed.), Williams and Wilkins Co., Baltimore. 2: 1105-1140.
- Ganesan, P. and Gnanamanickam, S.S. 1987. Biological control of *Sclerotium rolfsii* sacc. In peanut by inoculation with *Pseudomonas fluorescens*. *Soil Biol. Biochem.* 19(1): 35-38.
- Hema, C. Rao and Savalgi, V.P. 2017. Isolation and screening of nitrogen fixing endophytic bacterium *Gluconacetobacter diazotrophicus* GdS25. *Int. J. Curr. Microbiol. App. Sci.* 6(3): 1364-1373.
- Hu, X.F., Chen, J. and Guo, J.F. 2006. Two phosphate and potassium solubilizing bacteria isolated from Tiannu mountain, Zhejiang, China. *World J. Micro. Biotech.* 22: 983-990.
- Jadhav, R.N. 2013. Isolation of rhizobia from

- soybean cultivated in Latur area and study of its phosphate solubilizing ability. *Bioscience Discovery*. 4(1): 100-103.
- Parmar, K.B., Mehta, B.P. and Kunt, M.D. 2016. Isolation, characterization and identification of potassium solubilizing bacteria from rhizosphere soil of maize (*Zea mays*). *Int. J. Sci. Environ. Tech.* 5(5): 3030-3037.
- Qureshi, M.A., Shair, M.A., Iqbal, A., Ahtar, N. and Khan, A. 2011. Co-inoculation of phosphate solubilizing bacteria and rhizobia for improving growth and yield of mungbean (*Vigna radiata L.*). *J. Animal Plant Sci.* 21(3): 491-497.
- Rajendran, G., Singh, F., Desai, A. J. and Archana, G. 2008. Enhanced growth and nodulation of pigeonpea by co-inoculation of *Bacillus* strains along with *Rhizobium* sp. *Bioresource Tech.* 99: 4544-4550.
- Reis, V.M., Olivares, F.L. and Dobereiner, J. 1994. Improved methodology for isolation of *Acetobacter diazotrophicus* and confirmation of its habitat. *World J. Microbiol. Biotech.* 10: 101-104.
- Sagervanshi, A., Priyanka, K. and Anju, N. 2014. Media optimization for inorganic phosphate solubilizing bacteria isolated from agriculture soil. *Int. Life Sci. Pharma Res.* 2(3):245-255.
- Shete, M.H., Murumkar, D.R., Tirmali A.M. and Landge, K.B. 2019. Formulation of culture media for growth of nitrogen fixing, phosphate solubilizing and potash mobilizing bacteria in a consortium. *J. Pl. Dis. Sci.* 14(1): 41-46.
- Singh, A.K., Dipuraj, Harison, H., Yashab, K., Peter, J.K., Mishra, S.K. and Saxena, L. 2014. Optimization of production parameters and evaluation of shelf life of *Rhizobium* biofertilizers. *Elixir. Bio. Tech.* 67: 21787-21795.
- Sugumaran, P. and Janarthanam, B. 2007. Solubilization of potassium obtaining minerals by bacteria and their effect on plant growth. *World J. Agric. Sci.* 3(3): 350-355.
- Tarafder, H.K., Dey, A. and Dasgupta S. 2016. Co-inoculation of phosphate solubilizing bacteria and Rhizobia for improving growth and yield of mungbean (*Vigna radiata L.*). *Asian J. Soil Sci.* 11(1): 207-212.
- Sagervanshi, A., Priyanka, K. and Anju, N.

How to cite this article:

Ghadge, J. R., D. R. Murumkar and Shete, M. H. 2020. Development of Microbial Consortia of Nitrogen Fixing, Phosphate Solubilizing and Potash Mobilizing Bacteria for Optimizing Nutrient Supplementation to Soybean. *Int.J.Curr.Microbiol.App.Sci.* 9(07): 1889-1902.
doi: <https://doi.org/10.20546/ijcmas.2020.907.216>